

A Learning-Based Model for Imputing Missing Levels in Partial Conjoint Profiles

Eric T. Bradlow, Ye Hu, and Teck-Hua Ho*

January 26, 2004

* Eric T. Bradlow (email: ebradlow@wharton.upenn.edu) is an Associate Professor of Marketing and Statistics and Academic Director of the Wharton Small Business Development Center, The Wharton School of the University of Pennsylvania. Ye Hu (email: huye@wharton.upenn.edu) is a Visiting Assistant Professor, Krannert School of Management, Purdue University. Teck-Hua Ho (email: hoteck@haas.berkeley.edu) is William Halford Jr. Family Professor of Marketing, Haas School of Business, University of California, Berkeley. Professor Ho was partially supported by a grant from the Wharton-SMU Research Center, Singapore Management University. We thank David Robinson and Young Lee for their help in data collection and seminar participants at UC Berkeley, Singapore Management University, University of Michigan at Ann Arbor, the Marketing Science 2002 conference, and the BAMMCONF 2002 for their useful suggestions.

ABSTRACT

A Learning-Based Model for Imputing Missing Levels in Partial Conjoint Profiles

Respondents in a conjoint experiment sometimes are presented with successive partial product profiles. First, the authors model how respondents infer missing levels of product attributes in a partial conjoint profile by developing a learning-based imputation model that nests several extant models. The model is estimated using a hierarchical Bayesian approach to account for heterogeneity. The advantage of this approach over previous research is that it infers missing levels of an attribute not only from prior levels of the same attribute, but also from prior levels of other attributes, especially those that match the attribute levels of the current product profile. Second, the authors provide an empirical demonstration of their approach and test whether learning in conjoint studies occurs, to what extent, and in what manner it affects responses, partworths, and the relative importance of attributes. They show that the relative importance of attribute partworths can shift when subjects evaluate partial profiles, which suggests that consumers may construct rather than retrieve partworths and are sensitive to the order in which the profiles are presented. Finally, the results show that consumers' imputation processes can be influenced by manipulating their prior information about a product category. This research is of both theoretical and practical importance. Theoretically, it sheds light on how customers integrate different sources of information in evaluating products with incomplete attribute information. Practically, it highlights the potential pitfalls of imputing missing attribute levels using simple rules and a better behavioral model is developed for describing and predicting customers' ratings for partial conjoint profiles.

Keywords: Conjoint Analysis, Learning, Consumer Choice, Hierarchical Bayes Methods.

INTRODUCTION

Conjoint analysis is perhaps the most celebrated research tool in marketing. It has been applied to solve a wide variety of marketing problems ranging from understanding consumer preferences, estimating new product demand, to designing a brand new product line. The method involves presenting customers with a carefully chosen set of product profiles (called a test set) from the universal set (as defined by the levels of the attributes) and collecting their preferences, which could be ratings, rankings, or choices, for product profiles in the test set. The power of the method lies in its ability to extrapolate customers' preferences from this test set to the universal set. Clearly, the conjoint analysis method works better when the test set is small and the preference task difficulty is low. Both factors can be significantly influenced by the number of product attributes.

If the number of attributes is large (as in many high-tech durable products), a full-factorial experiment would require a respondent to assess their preferences for a large number of profiles, each consisting of many attributes. The large test set problem can be solved by using a fractional design (Plackett and Burman 1946; Green, Carroll, and Carmone 1978) that divides the test set among several respondents within a common customer segment.

There are two ways to solve the task difficulty problem. The first way is to use a self-explicated conjoint analysis (SECA) (Green 1984): consumers first rate the importance of the attributes, and then evaluate the attractiveness of each attribute level. By multiplying the normalized importance and attractiveness ratings, one can derive a consumer's overall preference for any profile. This approach typically requires the respondent to answer a smaller set of questions than a full profile judgment task, and avoids the complexity of judging a profile with too many attributes. However, the SECA method has its own problems, including the fact that attribute importance ratings by respondents are not always consistent with their preference decisions; the experimental condition of separating attribute and level ratings is artificial because real-life purchase decisions are made on whole products.

The second solution is to use orthogonal subsets of all the attributes (Green 1974), the so-called partial profile conjoint analysis (PPCA). Given that profiles with a smaller number of attributes may be

easier to rate (our experiment reveals that subjects took a significantly shorter time when they were asked to rate partial profiles), the PPCA approach decreases the difficulty of the rating task; however, at the same time, it may increase the number of profiles needed to determine consumers' utility functions. The PPCA approach typically assumes that the attributes that are missing do not impact the product evaluation. Several studies cast doubt on this assumption (e.g., Feldman and Lynch 1988; Huber and McCann 1982; Broniarczyk and Alba 1994). Consequently, standard rating conjoint analysis methods, applied to partial profiles, may not produce a highly predictive utility function. This paper investigates how subjects impute missing attribute levels when evaluating partial conjoint profiles. Our goals are to understand the dependency of ratings of current profiles on all available attribute information (in both the current and previously shown profiles), including a person's prior knowledge, and to provide insights as to how consumers may impute missing levels when evaluating partial conjoint profiles.

In this paper, we relax the "null effect for missing attribute" assumption and develop a probabilistic model of how respondents impute values for missing attributes based on a person's prior over the set of attribute levels, a given attribute's previously shown values, the previously shown values of the other attributes, and the covariation between attributes (both *a priori* and learned within the task itself). We conceptualize how consumers infer missing values via a pattern matching and learning process. Our model assumes that consumers learn and update after each stimulus (partial profile) about the pattern underlying the product attributes, their levels, and the correlations between them. How are strengths of patterns formed and updated? We assume that consumers have prior knowledge about the patterns and use knowledge about the product profiles acquired through the conjoint task to update their strengths. It is this dynamic process (learning about the attribute level occurrences and covariation between attributes) that we model and focus on in this paper.

We call the fundamental kernel of this updating structure a "pattern matching" learning model. That is, previously shown profiles that exhibit certain patterns among the attributes are used by the respondent to infer the missing attribute levels in the current profile. In essence, this approach can be viewed as a

time-varying, multi-way contingency table of latent counts for imputation (Little and Rubin 1987).

Consequently, the *order* in which profiles are presented matters in predicting preference ratings.

We model how people *rate* partial conjoint profiles over time. While rating-based methods may currently be less common than choice-based methods in practice (Wittink and Cattin 1989), our study is relevant for common applications of conjoint analysis in at least three ways¹:

1. Our research can influence the way that Adaptive Conjoint Analysis (ACA) is used in practice. The ACA engine (for example, equation (2) in ACA 5.0 Technical Report 2002) requires continuous strength of preference data, treated as a rating score, obtained for pairwise *partial* profiles. Although ACA can handle up to 30 attributes, it is suggested that each profile should contain no more than five attributes, a practice which has been brought into question by others (see for example, Green, Krieger, and Agarwal 1991; and subsequent response Johnson 1991). Our work is directly applicable to the ACA engine, which selects profile pairs based on utility balance, and, if those utilities are influenced by the missing attributes that do not cancel across choice pairs due to covariation, learning, etc., the resulting part-worths may be biased in some sense. Our model will help to quantify these biases or select pairs that indeed have the highest likelihood of canceling out on those missing attributes.
2. Our research has implications for the hybrid approach proposed by Srinivasan and Park (1997) and subsequently extended by Ter Hofstede et al. (2002). That is, both share the same data structure in which a subset of the most important attributes, based on self-explicated data, is used in a subsequent “full profile” conjoint study. The phrase full profile is in quotes because respondents are aware of all of the attributes before they rate the profiles that contain only a subset of attributes. In this manner, one could consider the profiles shown in their approaches “partial” and our model is able to shed light on the role of those attributes that are excluded in the rating task.
3. Our study is also relevant for choice-based conjoint (CBC) methods despite the assumption of ignorability across pairs of not shown attributes (Elrod, Louviere, and Davey 1992). Conceptually, all profiles, even if stated as full profile, have missing attributes that could be inferred. Therefore, despite

¹ We thank the editor and an anonymous reviewer for pointing out the link of our study to these approaches.

the common practice of stating “respondents were instructed that profiles were very similar in every respect except possibly for those attributes shown in the profile description”, it is an open empirical question whether respondents do, or possibly more importantly *can* follow this instruction. Hence our model can be used to check whether indeed subjects are able to ignore levels of those attributes that are not included in the study.

The rest of this paper is composed of four sections. In section 2, we develop our imputation model. In section 3, the design of our experiment is explained. In section 4, we estimate our model on two sets of experimental data and report the results. Section 5 contains conclusions, caveats, and areas for future research.

THE IMPUTATION MODEL

Notation and Model Set-up

We investigate how I individuals (indexed by $i = 1, \dots, I$) rate a series of product profiles (partial or full) in a conjoint experiment. Each product profile is characterized by J attributes (indexed by $j = 1, \dots, J$) and each attribute j has two levels. Each individual i rates T profiles (denoted by $M_i(t)$, $t = 1, \dots, T$) one-by-one. Individual i 's rating for product profile $M_i(t)$ is given by $y_i(t)$. Profile $M_i(t)$ takes a level of $x_{ij}(t) = 1$ or 0 for attribute j . We denote whether or not attribute j is missing in the t -th shown profile to respondent i by $r_{ij}(t)$. If attribute j is missing in profile $M_i(t)$, $r_{ij}(t)$ is zero, otherwise $r_{ij}(t)$ is 1. The basic premise of our model is that subject i does not ignore a missing attribute level but instead constructs an imputed value for it. Let $x'_{ij}(t)$ be that imputed value determined as follows:

$$x'_{ij}(t) = \begin{cases} x_{ij}(t) & \text{if } r_{ij}(t) = 1 \\ Ix_{ij}(t) & \text{if } r_{ij}(t) = 0 \end{cases}$$

Note that $Ix_{ij}(t)$ is a random variable, taking on the value of 1 or 0 (or in general the possible values of $x_{ij}(t)$), and the imputation modeling effort is to determine its probability distribution over the possible

levels. If an attribute is non-missing (i.e., $r_{ij}(t) = 1$), it is assumed that the shown attribute level, $x_{ij}(t)$, occurs with probability 1.

To determine the part-worths of the attributes, we postulate a regression with heterogeneous coefficients given by:

$$y_i(t) = \alpha_i + \sum_{j=1}^J \beta_{ij} x'_{ij}(t) + \varepsilon_i(t) \quad (1)$$

where β_{ij} is respondent i 's part-worth for attribute j .

Johnson, Levin, and their colleagues (Johnson and Levin 1985; Levin et al. 1986; Johnson 1987), however, suggest that subjects may have different part-worths for the same attribute when it is missing compared to when it is not. To control for this, the above regression equation is modified to yield:

$$y_i(t) = \alpha_i + \sum_{j=1}^J [\beta_{ij} x'_{ij}(t) + \beta'_{ij} r_{ij}(t)] + \varepsilon_i(t) \quad (2)$$

Table 1 shows the model's part-worths under different conditions. If subjects indeed have different part-worths when an attribute is missing, then β'_{ij} will be significantly different from zero.

[Insert Table 1 Here.]

Thus, our model nests the work of Johnson, Levin, and others and generalizes theirs by including imputed attribute levels $x'_{ij}(t)$ when $r_{ij}(t) = 0$.

Basic Ideas

Table 2 shows a hypothetical example that introduces the basic ideas of our imputation model, as well as demonstrates current extant models. There are four attributes (i.e., $J = 4$) and three profiles (i.e., $M_i(1)$, $M_i(2)$, $M_i(3)$). Each profile has one missing attribute (denoted as "MA") where subject i rates $M_i(1)$, $M_i(2)$, $M_i(3)$ sequentially. At time $t = 3$, we have $x_{i1}(3) = 1$, $x_{i3}(3) = 1$, $x_{i4}(3) = 0$. Attribute 2 is missing at time $t = 3$ and we denote its imputed level, 1 or 0, by $x'_{i2}(3)$. In a real experiment, the respondent is shown a

product profile at time 3 with only attributes 1, 3, and 4, and does not see “MA” for attribute 2; we include it in Table 2 only to describe the design.

[Insert Table 2 Here]

Assume that the subject has finished rating profiles $M_i(1)$ and $M_i(2)$, and profile $M_i(3)$ is the “current” product profile. To investigate how information from different attributes might influence the imputed value for attribute 2 at time $t = 3$, we divide the attributes into three types: 1) the omni-present (OM) set; 2) the presence-manipulated (PM) set that are present (non-missing PM); and (3) the presence-manipulated set that are missing (missing PM). The OM attributes are always presented while the PM attributes may or may not be. A PM attribute is called a “non-missing PM attribute” if it is *not* missing in the *current* conjoint profile. A PM attribute is called a “missing PM attribute” if it is missing in the *current* conjoint profile but may not be missing in others. In profile $M_i(3)$, attribute 1 is an OM attribute; attributes 3 and 4 are non-missing PM attributes; and attribute 2 is a missing PM attribute. Existing models utilize only the past information (values) from the currently missing PM attribute (attribute 2) for imputation of the missing level $x_{i2}(3)$. Our model uses all three sources, missing and non-missing PM attributes and OM attributes.

There are several different existing ways to treat missing attribute levels. The first way is to assume that respondents ignore them (Green 1974). Such an assumption implies that all the “MA”s in Table 1 are filled in as 0 (the default level). Formally, this assumption leads to the following prediction of the missing attribute level: $\Pr(x'_{i2}(3) = 0) = 1$ and $\Pr(x'_{i2}(3) = 1) = 0$. Note that, in this case, the imputation process of the missing levels depends on which level is coded as 0, a potential problem theoretically.

An alternative approach is to assume respondents impute values using all available information. That is, we assume that people infer the levels of the missing attributes from previously shown product profiles, and weight each profile “pattern” accordingly. This latter view is consistent with Meyer (1981) where he shows that when a subject has no information about certain attributes, that attribute is not ignored, but assigned a score equal to the individual’s adaptation level.

There are two common ways to model how consumers make inferences about missing attribute levels. One way is based on the so-called “recency effect” (Lynch and Srull 1982): people assume the missing attribute level to be the last level of the *same* attribute they saw. According to such a model, in Table 2, attribute 2 in profile $M_i(3)$ takes level 1 (following the level of attribute 2 in profile $M_i(2)$). Formally, we have: $\Pr(x'_{i2}(3) = 0) = 0$ and $\Pr(x'_{i2}(3) = 1) = 1$.

A second commonly used imputation approach is the averaging model (Yamagishi and Hill 1981); people impute the missing attribute level by averaging all the previously shown levels of the missing attribute. For example, in Table 2, this yields $\Pr(x'_{i2}(3) = 0) = 1/2$ and $\Pr(x'_{i2}(3) = 1) = 1/2$.

In imputing the missing values, the recency and averaging models make strong assumptions about the similarity between the current profile and the previous profiles. The recency-based model assumes that the current profile is “similar” only to the most recently shown profile and is dissimilar to the rest². The averaging model assumes that the current profile is equally similar to all the previously shown profiles. One would expect that some previously shown profiles are more similar (“count more” in imputing) to the current profile while others are less so.

All the above models utilize *only* past data from attribute 2 to impute $x'_{i2}(3)$. By doing so, they ignore two important pieces of information. First, there are the complete set of patterns shown to the subjects ($M_i(1), M_i(2)$), not just the values for attribute 2 ($x_{i2}(1), x_{i2}(2)$). Some of these patterns might occur more frequently, so their values for attribute 2 might be more salient and memorable. Second, levels of other attributes from the current profile (i.e., $M_i(3)$) might be diagnostic about the missing level. For example, if attribute 1 is negatively correlated with attribute 2, as in Table 2, a respondent might infer a 0 for Attribute 2 from a 1 in Attribute 1. Such a correlation structure could be based on people’s long-term memory or from the “learning” in the conjoint task. Huber and McCann (1982) showed that people use their belief of the correlation structure between price and quality to infer the missing price or quality when either one is missing. Broniarczyk and Alba (1994) also show that consumers’ intuitions (priors)

² Note that if $M_i(2)$ were to be first profile and $M_i(1)$ were to be the second profile, the prediction of the “recency model” would have been reversed (i.e., $\Pr(x'_{i2}(3) = 1)$). The averaging model, however, would have given the same prediction.

influence their inference making. Our model captures these covariances as well as the priors they “arrive” at the experiment with in a parsimonious way.

Like existing models, our imputation model derives probabilities that the missing PM attribute 2 in profile $M_i(3)$ takes a value of 1 or 0. The parameterization of the probabilities is based upon the work of Hoch, Bradlow, and Wansink (1999), which describes a similarity measure between a pair of categorical objects (conjoint profiles in this research), and Camerer and Ho (1998, 1999) and Ho and Chong (2003), which describe how learning and memory decay occur over time. The two basic concepts we utilize here are what we call “Pattern Matching” and “Experience Counts”. They are then combined to yield our imputation model which defines the probabilities over the missing attribute levels.

Three potential classes of models (Figures 1A-1C) are developed to demonstrate how the generality of our model is built up sequentially, using varying information sources.

[Insert Figures 1A, 1B, and 1C Here.]

The model in Figure 1A uses only previously shown information about the missing PM attribute (i.e., $j = 2$) to impute missing levels. It is a natural extension of the recency and averaging model that allows for decay. The imputation is based on historical levels of attribute 2 (0 in profile $M_i(1)$ and 1 in profile $M_i(2)$), but now with the more recent level (1 in profile $M_i(2)$) being potentially more influential. To capture the “recency effect” in a “decay-weighted” averaging model, we introduce a decay parameter, λ_2 ($0 < \lambda_2 \leq 1$, the subscript denoting attribute 2). We also introduce a new concept, “experience count (EC)” (Camerer and Ho 1999) such that $N_{ij}(t|l)$ denotes respondent i ’s latent EC of attribute j at time t , taking level l .

As illustrated in Figure 1A, $N_{i2}(3|0) = \lambda_2^2$ because $M_i(1)$ has an observed level of 0 on attribute 2 at time 1 (i.e., $x_{i2}(1) = 0$), and if $x'_{i2}(3)$ were to be imputed as a 0, λ_2 gets a power of 2 because there are two periods of time difference between profile $M_i(1)$ and profile $M_i(3)$ in which they would then match on PM attribute 2. Similarly, $N_{i2}(3|1) = \lambda_2$ because $M_i(2)$ has $x_{i2}(2) = 1$ and would match with $x'_{i2}(3)$ if it were to be imputed as a 1. Therefore, the probability that $x'_{i2}(3)$ takes level 1 or 0 is thus:

$$\Pr(x'_{i_2}(3) = 1) = \frac{N_{i_2}(3|1)}{N_{i_2}(3|1) + N_{i_2}(3|0)} = \frac{\lambda_2}{\lambda_2 + \lambda_2^2}$$

$$\Pr(x'_{i_2}(3) = 0) = 1 - \Pr(x'_{i_2}(3) = 1)$$

The averaging model is a special case of this class of models in which $\lambda_2 = 1$. The recency model has $\lambda_2 \rightarrow 0$, with $\Pr(x'_{i_2}(3) = 1) \rightarrow 1$ and $\Pr(x'_{i_2}(3) = 0) \rightarrow 0$.

The more general model in Figure 1B uses information from both the missing and non-missing PM attributes (attributes 3 and 4) for imputation. That is, in addition to using information from attribute 2 itself, we utilize possible conditional match patterns between profiles on the non-missing PM attributes. Since we assume that the missing attribute levels are not used for imputation³, we only need to check whether there is a match between $[x_{i_3}(3)]$ and $[x_{i_3}(2)]$ and between $[x_{i_4}(3)]$ and $[x_{i_4}(1)]$. Since $[x_{i_3}(3)]$ and $[x_{i_3}(2)]$ match, we expect $M_i(2)$ to influence imputation more than $M_i(1)$ on $M_i(3)$. Another decay parameter, λ_3 ($0 < \lambda_3 \leq 1$), is added to capture this reinforcement. Consequently, $N_{i_2}(3|1)$ becomes $\lambda_2 + \lambda_3$, whereas $N_{i_2}(3|0)$ stays the same. The probabilities that $x'_{i_2}(3)$ is imputed as 1 or 0 now becomes:

$$\Pr(x'_{i_2}(3) = 1) = \frac{N_{i_2}(3|1)}{N_{i_2}(3|1) + N_{i_2}(3|0)} = \frac{\lambda_2 + \lambda_3}{(\lambda_2 + \lambda_3) + \lambda_2^2}$$

$$\Pr(x'_{i_2}(3) = 0) = 1 - \Pr(x'_{i_2}(3) = 1)$$

Notice that $\Pr(x'_{i_2}(3) = 1)$ becomes larger, when compared to Figure 1A, as $M_i(2)$ (when compared to $M_i(1)$) is more similar to $M_i(3)$ than when we considered only the missing PM attribute.

The model in Figure 1C uses all of the available information from both the PM and OM attributes to impute the missing level. Following the procedure above, the ECs are $N_{i_2}(3|0) = \lambda_1^2 + \lambda_2^2$ and $N_{i_2}(3|1) = \lambda_2 + \lambda_3$. The corresponding probabilities now become:

$$\Pr(x'_{i_2}(3) = 1) = \frac{N_{i_2}(3|1)}{N_{i_2}(3|1) + N_{i_2}(3|0)} = \frac{\lambda_2 + \lambda_3}{(\lambda_2 + \lambda_3) + (\lambda_1^2 + \lambda_2^2)}$$

$$\Pr(x'_{i_2}(3) = 0) = 1 - \Pr(x'_{i_2}(3) = 1)$$

The most general model, Figure 1C, has two desirable properties:

³ This is an assumption/limitation of our approach that is discussed later and is an area for future research.

1. It uses all available information in the previously shown and current profiles in a sensible way.

Furthermore, the model highlights the potential pitfalls of the averaging and recency models. For example, it implies that these previous simpler models will yield the same prediction, given above, if $M_i(3)$ were to take any of these patterns $\{[1, MA, 1, 1], [1, MA, 1, 0], [1, MA, 0, 1], [1, MA, 0, 0], [0, MA, 1, 1], [0, MA, 1, 0], [0, MA, 0, 1], [0, MA, 0, 0]\}$, which seems very unlikely.

2. It allows respondents to apply different weights to different attributes depending on their preference.

General Formulation

In general, the pattern matching between two profiles can be formally defined as follows. Assume $r_{ij}(t) = 0$ and the level of attribute j at time t for respondent i is to be imputed. For each possible level of $x'_{ij}(t)$, we consider all previously *shown* profiles ($t' < t$) that have attribute j taking the same value (i.e., $x_{ij}(t') = x'_{ij}(t)$). That is, we find all t' in which the indicator function $I(x_{ij}(t'), x_{ij}(t)) = 1$. In addition, we set $I(x_{ij}(t), x_{ij}(t'))$ to 1 for those profiles $M_i(t')$ that have a match along a different attribute j' with the current profile $M_i(t)$, but also on the missing PM attribute j . We call this a conditional match-up model, as the pairs of profiles must match on the missing PM attribute for it to add to the EC.

It is also important to note the following “properties” of our pattern matching approach. (1) We do *not* match profiles based on imputed values of previous attributes, or in a more-than-one missing attribute case, imputed values of missing PM attribute(s) $j' \neq j$. (2) The way in which we “match” a given pattern is binary (yes it matched, no it did not). One could imagine a metric-based degree of matching model. We chose a binary Hamming metric approach as it is parsimonious, easy to describe, and is cognitively simple.

Let $N_{ij}(t|l_j)$ denote the latent EC of individual i for attribute j to take level l_j at time t . With attribute j as the missing PM attribute, our model for $N_{ij}(t|l_j)$ in complete generality is given by

$$N_{ij}(t|l_j) = N_{ij}(0|l_j) + \sum_{t'=1}^{t-1} \left\{ \lambda_{i(j,j)}^{t-t'} \cdot I(x'_{ij}(t), x_{ij}(t')) + \sum_{j'=1, j' \neq j}^J \left[\lambda_{i(j,j')}^{t-t'} \cdot I[x_{ij'}(t), x_{ij'}(t')] \cdot I[x'_{ij}(t), x_{ij}(t')] \right] \right\} \quad (3)$$

= “prior count” + {“missing PM count” + [“non-missing PM attribute + OM attribute count”]}

where $N_{ij}(0|l_j)$ denotes the “prior” count of person i on attribute j , level l_j at “time 0” and $0 < \lambda_{i(j,j)} \leq 1$ is the decay parameter for person i relating attribute j' ($j' = 1, \dots, J$) to j . $N_{ij}(0|l_j)$ allows for the possibility of prior knowledge of the marginal frequency of attribute levels and prior correlation between attribute levels.

In our experimental results, section 4, we fit a fairly general (reduced-form) version of the model (Equation (3)) that has the following set of specifications for λ_{ij} . This corresponds to Table 3, an example with digital cameras in which $j = \{1, 2, 3, 4\}$ are PM attributes (delay between shots, storage media, maximum resolution, and camera size) and $j = \{5, 6\}$ are OM attributes (price and mini-movie).

$$\begin{aligned} \lambda_{i(j,j')} &= \lambda_{ij} && \text{if } j' = j, j' \in PM, \forall j = 1,2,3,4 \\ \lambda_{i(j,j')} &= \lambda_{i75} && \text{if } j' \neq j, j' \in PM \\ \lambda_{i(j,j')} &= \lambda_{i6} && \text{if } j = \text{Maximum Resolution and } j' = \text{Price} \end{aligned}$$

Note that λ_{i6} is included in the model, as described in section 4, due to a prior manipulation of the covariance between price and maximum resolution.

[Insert Table 3 Here]

The structure described above defines the entire imputation process for partial profile conjoint designs as a time varying latent contingency table with counts, $N_{ij}(t|l_j)$, given in Equation (3). Hence, $\Pr(x'_{ij}(t) = l_j)$, $l_j = 1$ or 0, is given by

$$\Pr(x'_{ij}(t) = l_j) = \begin{cases} 1 & \text{if } r_{ij}(t) = 1 \text{ and } x'_{ij}(t) = l_j \\ 0 & \text{if } r_{ij}(t) = 1 \text{ and } x'_{ij}(t) \neq l_j \\ \frac{N_{ij}(t|l_j)}{N_{ij}(t|0) + N_{ij}(t|1)} & \text{if } r_{ij}(t) = 0 \end{cases} \quad (4)$$

That is, the probability a given attribute level is imputed, when that attribute is missing, is its “proportion” of the total EC for that attribute. As $N_{ij}(t|l_j)$ incorporates information across patterns to reinforce each pattern, and allows for differing importance across time, this model satisfies our basic pattern matching

and reinforcement requirements. When multiple attributes are missing, we assume independence of counts to derive the joint probability of the missing pattern; however, those counts that are correlated as attribute levels that occur together have counts that will be updated together, and prior counts that are related.

We denote the vector of imputed values at time t by a row vector $\mathbf{x}'_i(\mathbf{t}) = [x'_{i1}(t), x'_{i2}(t), \dots, x'_{ij}(t)]$. For a conjoint design with J attributes, where each attribute has two levels, the imputed value vector $\mathbf{x}'_i(\mathbf{t})$ may assume one of the $K = 2^J$ possible potential profiles. We denote these potential profiles by Z_k ($k = 1, \dots, K$) and we determine the probability that $\mathbf{x}'_i(\mathbf{t})$ equals potential profile Z_k as

$$\Pr(\mathbf{x}'_i(\mathbf{t}) = Z_k) = \prod_{j=1}^J \Pr(x'_{ij}(t) = l_j) \quad (5)$$

Heterogeneity

We allow for the rate of information decay for a specific attribute pattern to be individual and attribute specific, recognizing that considerable heterogeneity is likely to exist across persons in their decay attribute imputation parameters, λ_{im} ($m = 1, \dots, 6$). In addition, the basic parameters of the conjoint model, the individual conjoint intercepts, α_i , the attribute part-worths, β_{ij} and β'_{ij} , and the residual variances, may also contain considerable heterogeneity, yet share commonalities across the population of inference. To account for this heterogeneity in a coherent fashion, we nest our model in a Bayesian framework (Gelfand and Smith 1990). From equation (2), we have

$$\varepsilon_i(t) = y_i(t) - \left\{ \alpha_i + \sum_{j=1}^J [\beta_{ij} x'_{ij}(t) + \beta'_{ij} r_{ij}(t)] \right\}$$

We use an AR(1) (first-order autoregressive) model to capture potential correlation of error terms over time (that is, people may anchor somewhat on the previously provided rating):

$$\varepsilon_i(t) = \gamma_i \varepsilon_i(t-1) + u_i(t)$$

where

$$u_i(t) \sim N(0, \sigma_i^2) \quad (6)$$

In addition, we assume $\varepsilon_i(0) = 0, \forall i$.

Prior and hyperprior specifications for the conjoint parameters ($\forall i, j$) are given by⁴:

$$\begin{aligned} \gamma_i &\sim U(-1, 1) \\ \alpha_i &\sim N(\bar{\alpha}, \sigma_\alpha^2) \\ \beta_{ij} &\sim N(\bar{\beta}_j, \sigma_{j\beta}^2) \\ \beta'_{ij} &\sim N(\bar{\beta}'_j, \sigma_{j\beta'}^2) \\ \sigma_\alpha^2, \sigma_{j\beta}^2, \sigma_{j\beta'}^2 &\sim \text{Inv} - \Gamma(\cdot, \cdot) \end{aligned}$$

and prior specification for the attribute decay parameters, $0 < \lambda_{im} \leq 1$, given by

$$\lambda_{im} \sim \text{Beta}(a_m, b_m). \quad (7)$$

The prior ECs of each individual are assumed to follow a Poisson distribution with parameters varying by individuals and attribute levels:

$$N_{ij}(0 | l_j) \sim \text{Poisson}(\exp(\zeta_i + \omega_j)),$$

with slightly informative priors on ζ_i and ω_j . We note that $N(\mu, \sigma^2)$ denotes a normal distribution with mean μ and variance σ^2 , $U(g, h)$ a uniform distribution with lower bound g and upper bound h , $\text{Inv} - \Gamma(\cdot, \cdot)$ an inverse gamma distribution with corresponding parameters and $\text{Beta}(a, b)$ a beta distribution with parameters a, b . To complete the model specification, slightly informative hyperpriors were placed on σ_α^2 , $\sigma_{j\beta}^2$, and $\sigma_{j\beta'}^2, \forall j$ (inverse gamma distribution: $\text{Inv} - \Gamma(0.001, 0.001)$); $\bar{\beta}_j$ and $\bar{\beta}'_j, \forall j$ (normal distribution: $N(0, 0.001^{-1})$); and (a_m, b_m) (uniform distribution: $U(0, 1000)$). Numerous sensitivity analyses indicated that the results were not affected by the exact choice of uninformative hyperprior values.

To summarize, let the imputation model parameters be denoted by row vectors $\lambda_i = [\lambda_{i1}, \lambda_{i2}, \dots, \lambda_{i6}]$ (the length of λ_i varies with different models as described in section 4), and conjoint parameters by $\beta_i =$

⁴ We are aware that a more general setup would be to allow the β 's to follow a multivariate normal distribution with non-zero off-diagonal covariances. The current setup avoids over-parameterization of the model. It is commonly used in the economics literature, for example, Berry, Levinsohn, and Pakes (1995).

$[\beta_{i1}, \beta_{i2}, \dots, \beta_{i6}]$, and $\boldsymbol{\beta}'_i = [\beta'_{i1}, \beta'_{i2}, \dots, \beta'_{i6}]$. Given $\Pr(\mathbf{x}'_i(\mathbf{t}) = Z_k)$ from equation (5), the likelihood function is

$$L(\alpha_i, \boldsymbol{\beta}_i, \boldsymbol{\beta}'_i, \boldsymbol{\lambda}_i, \gamma_i, \sigma_i^2; \mathbf{y}_i) = \prod_{t=1}^{20} \left\{ \sum_{k=1}^{2^6} \left[\frac{1}{\sqrt{2\pi\sigma_i^2}} \exp\left(-\frac{([\varepsilon_i(t) | Z_k] - \gamma_i \varepsilon_i(t-1))^2}{2\sigma_i^2}\right) \cdot \Pr(\mathbf{x}'_i(\mathbf{t}) = Z_k) \right] \right\}. \quad (8)$$

That is, we integrate the conjoint regression model with respect to the imputation model, i.e. stick in the considered value for attribute j to person i for all possible profiles, and weight them by their probability of being the imputed corresponding level. We use the notation $[\varepsilon_i(t) | Z_k]$ to emphasize that the value of $\varepsilon_i(t)$ is conditional on profile Z_k . Also note that, although there are 2^6 potential profiles in our study, at each time t , only two profiles have non-zero $\Pr(\mathbf{x}'_i(\mathbf{t}) = Z_k)$ in the one-missing attribute case and only four in the two-missing attributes case.

Inferences from all models were derived by obtaining posterior samples using a Markov chain Monte Carlo sampler. All computation was performed using the software package WinBUGS, Bayesian Inference Using Gibbs Sampling (<http://www.mrc-bsu.cam.ac.uk/bugs/>)⁵. All of the results reported in section 4 are the posterior means obtained from aggregating the draws of three runs of the sampler from different starting points with a burn-in period of 6,000 draws, and a total run length of 10,000 draws. Convergence was assessed using the F-test approach of Gelman and Rubin (1992).

Due to the complexity of the model, the fact that many readers may lack familiarity with the WinBUGS software program, and a desire for other researchers to easily apply our model, we have included an annotated version of the WinBUGS code for our most general model in an online appendix at http://mktgweb.wharton.upenn.edu/ebradlow/research_files.htm.

⁵ To assess the ability of our most general model (Model 6) in recovering the true underlying model structure, we ran a simulation study using synthetic data. The simulation results indicate that the model is able to recover the underlying conjoint regression coefficients (α_i , β_{ij} , β'_{ij} , and γ_i) very accurately and the imputation parameters (λ 's) with reasonable accuracy. Details are available from the authors upon request.

EXPERIMENT

An experiment was designed to provide a basic demonstration of our model on rating conjoint data with missing attributes. Our interest lies in providing a demonstration of our approach, as well as to begin a preliminary understanding of:

1. Do people use missing attributes, and their levels, to evaluate products?
2. If yes, do they infer missing attribute levels from all of the information they learn about the product profiles; do they reinforce patterns?

We assume that a consumer has minimal prior information about the product, although we do estimate this as given in Equation (8). Therefore, we are able to impose a “prior” structure that varied across respondents in a systematic way (described below). First, through a learning process, we create a “prior” for each individual by controlling the products that she sees in a “learning phase”. Second, we ask participants to rate products with (or without, in the control group) missing attributes. A second control group worked on a self-explicated conjoint task to act as a second baseline.

Stimulus

Digital cameras were selected for this experiment as we wanted a relatively new product category where the frequency of attribute levels and the correlation structure of the attributes are mostly unknown. This would allow us to manipulate the frequencies and impose a prior as desired. From our demographic questions, less than 10% of our subjects owned digital cameras or claimed to have extensive prior expertise. We chose digital cameras with 6 attributes (as the full profile task) as research that we conducted indicated that digital cameras could be described well using 6 features. A summary of the digital camera attributes used are listed in Table 3. In our experimental condition, all attributes are simplified to have two realistic levels. This product set-up provides a stylized empirical test of our model.

Experimental Groups

The experiment was designed to run on a university network. One hundred and thirty undergraduate students from a large east-coast university participated in the experiment. Subjects were obtained and assigned to treatments by taking six sections of a large class and randomly assigning one section each to receive the self-explicated and full profile (0 missing) cases, and two sections each to receive the one missing and two missing attribute cases. This resulted in four groups with group sizes 17, 23, 47, and 43, respectively⁶. Group size differences were due to different section sizes and the participation rate of students in those sections. Across the conditions, less than 10% owned digital cameras; 40% are females and 60% are males.

The experiment was composed of two phases: the learning (prior) phase and the rating phase. In the learning phase, the subjects were provided with text information about digital cameras and their attributes. They were then shown 20 digital camera profiles listed in a single table. We control the consumers' priors by manipulating the digital camera profiles they see in the learning phase. In the rating session, the subjects were asked to rate, on a 0-9 Likert scale, the attractiveness of different digital cameras (some with partial product profiles depending on the treatment condition).

Learning Phase

In the learning phase, all subjects were shown 20 digital camera profiles. The priors of the subjects before the rating phase were manipulated by the learning phase profiles. The purpose of this learning phase manipulation is two-fold. First, if the relationship, say between price and maximum resolution, as described next, can be influenced by showing subjects profiles of a given structure, then managerial practice would suggest prior manipulations of this type could be valuable. Secondly, we wanted to test out our model, for a given attribute correlation. As we manipulated the priors between digital camera

⁶ We note that a better design would have been to randomly assign people and not sections. Study 2 utilized random assignment.

price and *maximum resolution*, we wanted to test if λ_{i6} would impact the EC for resolution when it is missing (price is never missing as it is an OM attribute).

Each subject was randomly assigned to one of eleven prior coincidence structures representing a different level of coincidence between price and maximum resolution (while keeping the coincidences between other attributes orthogonal.) Specifically, they were assigned to read a table with a specific coincidence value (between 0 and 10) between price and maximum resolution. For example, a coincidence value of 10 indicates that among the 10 profiles that have low price (\$159), all have low resolution (800X600); a value of zero would indicate that among the 10 profiles that have low price (\$159), all of them have high resolution (1024X768). Such coincidence structure could potentially affect $\Pr(x'_{ij}(t) = l_j)$ if the learning phase carries over to the rating phase. That is, we will test empirically the ability to manipulate the rating phase data by co-varying price and maximum resolution at levels 0, 1, ..., 10 in the learning phase, and then estimating λ_{i6} in our model and seeing its correlation with the subject's prior manipulation.

To ensure that subjects followed and attended to all information in the table of 20 learning profiles provided, they were asked to count five of the pairwise coincidences after they had read the tables. Among the five questions, one of them asked the subjects to count the coincidence between price \$159 and maximum resolution 800X600, the manipulated coincidence, while the other four questions were randomly chosen to ask the subjects to count other coincidences. The sequences of these questions were randomized so as not to bias the results. Their responses to these questions suggested that they had paid attention to the coincidence counts.

Rating Phase

In the rating phase, the design is orthogonal. In the one or two missing attribute group, one or two out of *four* PM attributes are removed from the designed conjoint cards, respectively. We fixed two attributes to be OM as we wanted to see the impact of imputation of missing levels on observed attribute part-worths. We utilized a Plackett-Burman Design (Plackett and Burman 1946; Green, Carroll, and Carmone

1978) to create the profile cards. The sequences in which the profile cards were shown were generated randomly and varied across respondent. Each subject saw 24 profiles in the rating phase. Debriefing questions after the experiments provided evidence that the subjects do “notice” that attributes are missing and utilize this fact in their ratings. Response time was recorded, which could be used as a proxy for the difficulty of the task. Each response time was the time (in seconds) measured between successive rating score responses being keyed in. An analysis of the response time data across missing attribute conditions (0, 1, and 2) indicated that respondents in the two missing case, spent considerably less time than in either of the other two cases ($p < 0.01$), corresponding to an average of 31 seconds less across the 24 profile rating tasks. No significant differences were found in response time between the zero and one missing attribute conditions.

RESULTS

We utilized the first 20 profiles for each subject to the calibrate model, and the last four as holdout for validation.

We estimated a total of six models, with differing degrees of generality. These models are grouped into four categories: (1) prior models, (2) imputation based on missing PM attributes only, (3) imputation based on missing and non-missing PM attributes, and (4) imputation based on the OM attribute (Price), missing and non-missing PM attributes. Table 4 shows these models and their relationships for both the one and two missing attribute cases. The estimation was done using the Bayesian hierarchical structure described in the “Heterogeneity” section.

[Insert Table 4 Here]

Table 5 shows the relative performance for the six models. We show results for the three extant models (ignore, recency, and averaging), as well as one model in each of the three classes (Models 4, 5, and 6). For each model, we report (1) the log of the marginal likelihood as computed by the log of the harmonic mean of the likelihood values (Congdon 2001, pp. 475), and (2) mean absolute errors (MAE),

both in sample and out of sample. Using all three measures, our models (Models 4-6) perform better than the prior models (Models 1-3). Specifically, Models 4 through 6 consistently perform better as more information is used for imputation.

[Insert Table 5 Here]

Imputation Based on Missing PM Attributes

As discussed above, the extant models assume that subjects either ignore the missing attribute, use the most recent occurrence of a missing attribute, or compute the average of all past occurrences. Out of models 1-3, the averaging model (Model 3) performs better in terms of log-marginal likelihood and in- and out-of-sample mean absolute error (MAE). Model 4 relaxes the assumptions of Model 3: it allows a separate λ for each missing PM attribute, which decays geometrically.

Compared to the averaging model, Model 4 performs better in terms of log-marginal likelihood, in sample and out of sample MAE. Such results suggest that the relaxation of allowing for heterogeneous geometric decay helps in terms of model performance and better captures the actual rating process when missing attributes exist.

Imputation Based on Missing and Non-missing PM Attributes

The above class of model assumes that subjects impute a missing level of an attribute using only information within that PM attribute. A natural extension is to account for covariation from the non-missing PM attributes. In Model 5, we assume each attribute to take a different λ when it is present and when it is missing; however, the value of λ is assumed to be common across all PM attributes when they are non-missing. As indicated in Table 5, Model 5 fits better than Models 1-4 in terms of log-marginal likelihoods, in-sample MAE, and out-of-sample MAE.

Imputation Based on OM Attribute, Missing, and Non-Missing PM Attributes

To fully test the information used by the subjects when inferring missing attribute levels, in addition to the last set of models, we added price (an OM attribute) to impute the missing level of maximum

resolution. Recall that we manipulated the correlation structure between price and resolution in the learning phase. Model 6 extends Model 5 by allowing price to be used in the imputation process for missing maximum resolution levels. Such a relaxation improves the log harmonic mean likelihood, and the in-sample and out-of-sample MAE. The results suggest that subjects do use OM attributes to infer missing attribute levels; albeit whether this goes beyond price, to other product domains, etc., is an open question. In fact, Model 6 outperforms Models 1-3 significantly by all the measures we considered in Table 5. Specifically, Model 6 decreases the out-of-sample MAE by 4.1%, 5.3%, and 6.5% over Models 1-3, respectively, in the one-missing attribute case; and 1.6%, 3.0%, and 5.9% over Models 1-3, respectively, in the two-missing attributes case.

A more detailed analysis at the individual-level between the estimated effect λ_{i6} (Price and Maximum Resolution) and the prior manipulated covariation between price and maximum resolution (0, 1, ..., 10) was done in a number of ways. (1) First, we note that λ_{i6} is significantly different from zero (the [2.5%, 97.5%] percentile of its posterior is [0.151, 0.315] in the one-missing attribute case and [0.628, 0.863] in the two-missing attributes case), suggesting significant effects overall. (2) An analysis at the individual-level (without shrinkage), indicated a significant effect in the one-missing attribute case (correlation = 0.345, $p = 0.017$) and insignificant in the two-missing case, between the prior manipulation and λ_{i6} . Overall, these findings suggest that the subjects' "priors" could be manipulated to influence the way they infer missing attributes, but to what extent remains an open empirical question.

The average λ values and their standard deviations of the best-fitting model (Model 6) are provided in Table 6. The estimated λ values are different in the one- and two-missing attributes cases, which is not surprising: when different number of attributes are missing, the weights which reflect how information from non-missing attributes is used change accordingly.

[Insert Table 6 Here]

Notice all the λ values are significantly larger than 0 and smaller than 1, indicating the actual imputation procedure is different from the pure effect of any one of Models 1-3.

Estimated Part-worths and Priors

Table 7 reports the mean and standard deviation of the part-worths of Model 6, the best fitting model. β'_{ij} , ($j = 1, \dots, 4, \forall i$) are the part-worths of the attributes when they take level 0 and are present (compared to taking level 0 and not being present, i.e. being imputed as a zero). $\beta_{ij} + \beta'_{ij}$, ($j = 1, \dots, 4, \forall i$) are the part-worths of the attributes when they take level 1 and are present (compared to taking level 0 and not being presented). To compare our results, therefore, with traditional conjoint part-worths, we note that with *all attributes present* and the “low” level attributes coded as zero (as is standard), the part-worths represent the difference in utility between the high and low attribute levels. To align with our case, the “traditional part-worths” from our model are β_{ij} , ($j = 1, \dots, 4, \forall i$); that is, the effect of being high when shown minus the effect of being low when shown.

As mentioned earlier, we get a “bonus” in that we can assess the effect of imputed versus not imputed attribute levels in our conjoint design, in addition to level 1 (high) versus level 0 (low). In our model, these are the part-worths, β'_{ij} . We find that all elements β'_{ij} ($j = 1, \dots, 4, \forall i$) have 95% posterior interval that do not contain 0, which means that when an attribute is present, it is given significantly greater weight. This finding is consistent with extant research (Meyer 1981; Levin et al. 1986; Johnson 1987; Louviere and Johnson 1990). Alternatively, we note that the combined tests (present or not, and levels 1 or 0), for each of the attributes, suggest that it is not the attribute level inferred, but the presence of the attribute that influences the weight put on the attribute. We believe this is an interesting area for future study.

Table 8 reports the relative importance of (traditional) part-worths, i.e. when high level is shown versus low level is shown, from Model 6, and from the case where there are no missing attributes. While we observe relatively high stability in the rankings (for instance, storage and size are always last, resolution is always most important, and the other three are relatively close in importance), there are changes in the magnitude of the relative importance of the part-worths. This finding, that part-worths themselves are “biased” (as compared to the full profile condition), is consistent with extant research

(Levin et al. 1986; Johnson 1987; Louviere and Johnson 1990). However, given that we also find that the relative rankings stay fairly stable, there is *prima facie* evidence that similar rating processes are going on. Interestingly, in the two-missing attributes case, when less attribute levels are available for imputation, the more important attributes in the no-missing case become less important, the less important ones become more important. Thus, there is a “regression” effect in part-worths when subjects evaluate partial profiles when less information is provided.

We note, that one way to interpret the observed changes in part-worths is that consumers “construct” rather than “retrieve” utilities. Since the set of all available information changes with successive profiles, the utilities can change even for identical profiles if they appear at different points in time. This view is not new and has been established by consumer researchers (Bettman and Zins 1977; Payne, Bettman, and Johnson 1992).

[Insert Table 7 Here]

[Insert Table 8 Here]

Finally, we report on the model results with regards to the carry-over effect from one rating’s error $\varepsilon_i(t-1)$ to another, and from the priors $N_{ij}(0|l_j)$. The AR(1) carry-over effect is statistically significant with mean of $\gamma \approx 0.1$ (in both one- and two-missing attributes cases), suggesting that people do anchor somewhat on previous values. This result suggests that the order in which previous profiles are presented could influence the subjects’ rating for a current profile.

None of the estimated prior parameters ζ_i ($i = 1, \dots, 47$) and ω_j ($j = 1, \dots, 6$) is significantly different from zero according to the [2.5%, 97.5%] percentile of their posterior draws in the one-missing attribute case, an indication that the subjects have a weak prior on the product category. Consequently, the average values of the experience counts, as shown in Table 9, for Model 6, are typically small. These initial experience counts thus exert some minor influence on the imputation of the early profiles but decay quickly when more profiles are shown. However, some of the prior parameters become significant in the two-missing attributes case. Specifically, ω ’s for resolution and size, the PM attributes people are

probably most familiar with, are fairly significant. This shows that when less information becomes available, people may depend more on their “priors” to make judgments. This certainly requires further study beyond the empirical example provided here.

[Insert Table 9 Here]

Robustness of Results

In our experiment, we “imposed” a prior on subjects’ beliefs about the relationship between price and maximum resolution via the learning phase, and subsequently measured whether it “held up” in the calibration phase. One may wonder whether our “process” of having people count relationships between pairs of attributes (that would normally not be done in practice), could bias people towards imputing attribute levels when they are missing due to priming⁷. To check whether our results are robust to this manipulation, we ran a second study, with a 0-missing and 1-missing attribute case only, that did not include a learning phase; yet in all other ways was identical to the first study. Our goal was to demonstrate the existence of imputation (as in our earlier experiment), and to replicate the patterns of superiority in for Models 4-6 over Models 1-3 (the learning-based models defeat the simpler models).

Specifically, ninety-one subjects from a large West Coast university, to partially fulfill requirements for a course, were obtained for our conjoint computer-based study of digital cameras with the same 6 attributes as in our Study 1. Subjects were randomly assigned to either the zero-missing case as a baseline (41 subjects), or the one-missing case (50 subjects). As in Study 1, the first twenty rating tasks were utilized for calibration of the model, and the remaining four for out-of-sample validation. Profiles were presented in a random order within each design.

[Insert Tables 10, 11, and 12 Here]

A detailed set of findings for this study are in Tables 10, 11, and 12, but at a summary level, our findings are as follows. First, an identical pattern of overall fit, both in and out of sample is found as in Study 1, in that the recency model has the worst fit, followed by the model which ignores the missing

⁷ We would like to thank the editor and an anonymous reviewer for suggesting this and the second study.

attributes and the averaging model, and then the three learning-based models. Other findings, as indicated by the mean value of $\gamma = 0.167$, and the pattern of relative part-worths for Model 6 (the best fitting model) indicate that our findings are robust overall to the learning phase manipulation, and were replicated.

CONCLUSIONS AND FUTURE RESEARCH

We have developed a learning model to describe how consumers impute missing levels in partial conjoint profiles. In our model, consumers match patterns and develop inferences based on their prior exposures. Our model extends the Averaging and Recency models and shows that consumers may infer missing attribute levels using both missing and non-missing attribute information. We have shown that our best-fitting models outperform the prior models both in-sample and out-of-sample.

The Ignore model is inadequate because consumers appear to consider missing attribute levels. Neither the Averaging nor Recency models does significantly better because consumers impute missing attribute levels using prior levels of non-missing attributes (whether PM or OM). At the same time, significant correlation between the manipulated coincidence in the learning phase and the estimated decay parameter provides evidence that the consumer's prior could be influenced by "communication" and "experience". Consequently, managers may be able to influence the overall attractiveness of a product to a consumer by making the consumers learn "prior knowledge" that favors the product.

This research has two caveats. First, the product used in our experiment has only six attributes, each with two levels. Thus, our study is best considered as a demonstration of the potential of our imputation model for predicting preferences in more complicated product categories. Second, our rating-based conjoint experiment does not allow us to provide direct evidence of the applicability of our model to CBC, although theoretically, such application is possible as described above.

We see at least three future research opportunities:

1. An interesting area to pursue is to model the trade-off between the number of profiles shown and the number of attributes shown in each profile. From an econometric perspective, it would be

interesting to keep the total number of attribute levels shown fixed and see how different combinations of the number of attributes and profiles would lead to different levels of information content.

2. As indicated in Section 2, we assume here a pattern matching model in which attributes either match or do not (0/1). A more general distance model can explicitly account for the relative differences between attribute levels. Such machinery is already in the marketers' toolbox as MDS (Multiple Dimensional Scaling) studies are used for such purposes. Thus, two promising areas for future studies would be: (a) to combine conjoint analysis and MDS studies to impute missing attribute levels, and (b) to create a "latent" perceptual mapping model for missing attribute levels in conjoint.
3. As mentioned previously, and shown in earlier research work, the missingness of attributes may indeed change the relative importance of attributes. While our work confirms that, and for the most important attributes, whether this is true generally is unclear, and what may moderate this effect may also be of interest. Thus, it would be interesting to conduct future studies to determine the degree of this change and its moderating variables.

In conclusion, we believe that the general theoretical framework presented here, and its empirical validations, are a good first step which we hope will lead to a future stream of managerially important research.

REFERENCES

- ACA 5.0 Technical Paper, Sawtooth Software Technical Paper Series (2002),
www.sawtoothsoftware.com.
- Berry S., J. Levinsohn, and A. Pakes (1995), "Automobile Prices in Market Equilibrium," *Econometrica*, 63(4), 841-90.
- Bettman, J. R., and M. Zins (1977), "Constructive Processes in Consumer Choice," *Journal of Consumer Research*, 4(2), 75-85.
- Broniarczyk, S. M. and J. W. Alba (1994), "The Role of Consumers' Intuitions in Inference Making," *Journal of Consumer Research*, 21(3), 393-407.
- Camerer, C. and T.-H. Ho (1998), "EWA Learning in Coordination Games: Probability Rules, Heterogeneity, and Time Variation," *Journal of Mathematical Psychology*, 42(2-3), 305-26.
- and —— (1999), "Experience-Weighted Attraction Learning in Normal Form Games," *Econometrica*, 67(4), 827-74.
- Congdon, P. (2001), *Bayesian Statistical Modeling*, West Sussex, England: John Wiley & Sons, Inc.
- Elrod, T., J. J. Louviere, and K. S. Davey (1992), "An Empirical Comparison of Ratings-Based and Choice-Based Conjoint Models," *Journal of Marketing Research*, 29(3), 368-77.
- Feldman, J. M. and J. G. Lynch, Jr. (1988), "Self-generated Validity and Other Effects of Measurement on Belief, Attitude, Intention, and Behavior," *Journal of Applied Psychology*, 73(3), 421-35.
- Gelfand, A. E., and A. F. M. Smith (1990), "Sampling-Based Approaches to Computing Marginal Densities," *Journal of the American Statistical Association*, 85(410), 398-409.

Gelman, A., and D. B. Rubin (1992), "Inference from Iterative Simulation Using Multiple Sequences," *Statistical Science*, 7(4), 457-511.

Green, P. E. (1974), "On the Design of Choice Experiments Involving Multifactor Alternatives," *Journal of Consumer Research*, 1(2), 61-8.

—— (1984), "Hybrid Model for Conjoint Analysis: An Expository Review," *Journal of Marketing Research*, 21(2), 155-69.

Green, P. E., J. D. Carroll, and F.J. Carmone (1978), "Some New Types of Fractional Factorial Designs for Marketing Experiments," In J.N. Sheth (Ed.), *Research in Marketing* 1, 99-122. Greenwich: JAI.

Green, P. E., A. M. Krieger, and M. K. Agarwal (1991), "Adaptive Conjoint Analysis: Some Caveats and Suggestions," *Journal of Marketing Research*, 28(2), 215-22.

Ho, T.-H. and J.-K. Chong (2003) "A Parsimonious Model of SKU Choice," *Journal of Marketing Research*, 40 (3), 351-65.

Hoch, S. J., E. T. Bradlow, and B. Wansink (1999), "The Variety of an Assortment," *Marketing Science*, 18(4), 527-46.

Huber, J. and J. W. McCann (1982), "The Impact of Inferential Beliefs on Product Evaluations," *Journal of Marketing Research*, 19(3), 324-33.

Johnson, R. D. (1987), "Making Judgments When Information is Missing: Inferences, Biases, and Framing Effects," *Acta Psychologica*, 66(1), 69-82.

Johnson, R. M. (1991), "Comment on 'Adaptive Conjoint Analysis: Some Caveats and Suggestions'," *Journal of Marketing Research*, 28(2), 223-5.

Johnson, R. D. and I. P. Levin (1985), "More than Meets the Eye: The Effect of Missing Information on Purchase Evaluations," *Journal of Consumer Research*, 12(2), 169-77.

Levin, I. P., R. D. Johnson, P. J. Deldin, L. M. Carstens, L. J. Cressey, and C. R. Davis (1986), "Framing Effects in Decisions with Completely and Incompletely Described Alternatives," *Organizational Behavior and Human Decision Processes*, 38(1), 48-65.

Little, R. J. A. and D. B. Rubin (1987), *Statistical Analysis with Missing Data*, New York: John Wiley & Sons, Inc.

Louviere, J. J. and R. D. Johnson (1990), "Reliability and Validity of the Brand-Anchored Conjoint Approach to Measuring Retailer Images," *Journal of Retailing*, 66(4), 359-81.

Lynch, J. G., Jr. and T. K. Srull (1982), "Memory and Attentional Factors in Consumer Choice: Concepts and Research Methods," *Journal of Consumer Research*, 9(1), 18-37.

Meyer, R. (1981), "A Model of Multiattribute Judgments Under Attribute Uncertainty and Informational Constraint," *Journal of Consumer Research*, 18(4), 428-41.

Payne, J. W., J. R. Bettman, and E. J. Johnson (1992), "Behavioral Decision Research: A Constructive Processing Perspective," *Annual Review of Psychology*, 43, 87-131.

Plackett, R. L. and J. P. Burman (1946), "The Design of Optimum Multifactorial Experiments," *Biometrika*, 33(4), 305-25.

Spiegelhalter, D.J., A. Thomas, N. G. Best, and W. R. Gilks (2000), *WinBUGS (Version 1.3)* [Computer software], Cambridge, UK: MRC Biostatistics Unit, Institute of Public Health.

Srinivasan, V., and C. S. Park (1997), "Surprising Robustness of the Self-Explicated Approach to Customer Preference Structure Measurement," *Journal of Marketing Research*, 34(2), 286-91.

Ter Hofstede, F., Y. Kim, and M. Wedel (2002), "Bayesian Prediction in Hybrid Conjoint Analysis," *Journal of Marketing Research* 39(2), 253-61.

Wittink, D. R., and Cattin, P. (1989), "Commercial Use of Conjoint Analysis: An Update," *Journal of Marketing*, 53(3), 91-6.

Yamagishi, T., and C. T. Hill (1981), "Adding versus Averaging Models Revisited: A test of a Path-Analytic Integration Model," *Journal of Personality and Social Psychology*, 41(1), 13-25.

Table 1. Part-worths of the same attribute

		Attribute Level	
		High: $x'_{ij}(t) = 1$	Low: $x'_{ij}(t) = 0$
Attribute Shown?	Yes	$\beta_{ij} + \beta'_{ij}$	β'_{ij}
	No	β_{ij}	0

Table 2. An illustrative example

Time (t)	OM	PM		
	Attribute 1	Attribute 2	Attribute 3	Attribute 4
1	1	0	MA*	1
2	0	1	1	MA
3	1	MA	1	0

*MA: missing attribute

Table 3. Digital camera attributes

Attribute	Delay Between Shots	Storage Media	Maximum Resolution	Camera Size	Price	Mini-Movie
Abbreviation	Delay	Storage	Resolution	Size	Price	Mini-Movie
Level 0	4 Seconds	Floppy Disk	800X600	SLR	\$239	No
Level 1	2 Seconds	Removable Memory	1024X768	Medium	\$159	Yes

Table 4. Description of models

Model Category	Model	Number of individual level parameters	λ Values of PM Attributes				Non-Missing PM	λ Values of OM Attributes	
			Missing PM					Price	Mini-Movie
			Delay	Storage	Resolution	Size			
Prior Models	1 §	12	---				---	---	---
	2 †	12	$\lambda_1 = \lambda_2 = \lambda_3 = \lambda_4, \rightarrow 0$				---	---	---
	3 ‡	12	1				---	---	---
Imputed based on missing PM	4	14	λ_1	λ_2	λ_3	λ_4	---	---	---
Imputed based on missing and non-missing PM	5	17	λ_1	λ_2	λ_3	λ_4	λ_5	---	---
Imputed based on OM & missing and non-missing PM	6	18	λ_1	λ_2	λ_3	λ_4	λ_5	λ_6	---

§ Ignore-missing model

† Recency Model

‡ Averaging Model

Table 5. Performance of different models

Model	1-missing			2-missing		
	Log- Harmonic Mean of Likelihood	Mean absolute error		Log- Harmonic Mean of Likelihood	Mean absolute error	
		In-sample	Out-of-sample		In-sample	Out-of-sample
1	-1285	0.807	1.366	-1232	0.861	1.382
2	-1245	0.778	1.386	-1187	0.823	1.390
3	-977	0.733	1.314	-685	0.500	1.381
4	-904	0.694	1.310	-580	0.461	1.360
5	-884	0.602	1.292	-513	0.459	1.341
6	-782	0.580	1.281	-356	0.434	1.300

Table 6. λ 's of the best-fitting model (Model 6)

Model	Values	Missing PM				Non-missing PM	OM (price)
		Delay	Storage	Resolution	Size		
1-missing	Average	0.306	0.095	0.829	0.240	0.977	0.247
	SD	0.056	0.066	0.044	0.067	0.018	0.046
2-missing	Average	0.981	0.429	0.032	0.070	0.697	0.743
	SD	0.012	0.091	0.018	0.061	0.074	0.070

Table 7. Average and standard deviation of the part-worths of the best-fitting model (Model 6)

Model	Coefficients	Delay	Storage	Resolution	Size	Price	Mini-Movie	Intercept	
1-missing	β_{ij}	Mean	1.328	0.046	1.384	0.092	0.829	1.296	0.592 (0.426)
		SD	0.124	0.182	0.133	0.144	0.124	0.158	
	β'_{ij}	Mean	0.844	0.745	0.876	0.737	---	---	
		SD	0.117	0.145	0.124	0.136	---	---	
2-missing	β_{ij}	Mean	1.001	0.565	1.267	0.469	1.000	1.085	0.526 (0.140)
		SD	0.004	0.177	0.088	0.150	0.004	0.130	
	β'_{ij}	Mean	1.196	1.238	1.168	1.119	---	---	
		SD	0.065	0.106	0.063	0.063	---	---	

Table 8. Comparison of relative importance of part-worths

Model	Delay	Storage	Resolution	Size	Price	Mini-Movie
0-missing case	0.126	0.029	0.317	0.056	0.198	0.275
1-missing case, Model 6 (β_{ij})	0.267	0.009	0.278	0.018	0.167	0.261
2-missing case, Model 6 (β_{ij})	0.186	0.105	0.235	0.087	0.186	0.201

Table 9. Average of estimated priors $N_j(0|\cdot)$

Models	1-missing		2-missing	
	0	1	0	1
Delay	1.025	0.931	4.198	2.461
Storage	0.194	0.722	2.898	4.812
Resolution	0.422	0.539	10.068	16.816
Size	0.453	0.441	4.439	4.858
Price	0.967	1.004	2.062	2.006
Mini-Movie	1.005	1.017	2.002	2.022

Table 10. Performance of different models (no learning)

Model	1-missing		
	Log- Harmonic Mean of Likelihood	Mean absolute error	
		In-sample	Out-of-sample
1	-582	0.905	1.182
2	-599	0.928	1.214
3	-526	0.866	1.182
4	-459	0.820	1.177
5	-424	0.808	1.167
6	-414	0.802	1.159

Table 11. Comparison of relative importance of part-worths (no learning)

Model	Delay	Storage	Resolution	Size	Price	Mini-Movie
0-missing case	0.144	0.114	0.308	0.026	0.190	0.217
1-missing case, Model 6 (β_j)	0.162	0.114	0.254	0.028	0.204	0.237

Table 12. Average of λ 's of the best-fitting model (Model 6, no learning)

Model	Values	Missing PM				Non-missing PM	OM (price)
		Delay	Storage	Resolution	Size		
1-missing	Average	0.624	0.126	0.555	0.037	0.065	0.598
	SD	0.092	0.037	0.068	0.013	0.027	0.128

Figure 1A. Imputing from missing PM only




Figure 1B. Imputing from missing and non-missing PM




Figure 1C. Imputing from OM, missing and non-missing PM

