

Lecture 8

Models for Censored and Truncated Data - Tobit Model

1

Censored and Truncated Data

- In some data sets we do not observe values above or below a certain magnitude, due to a censoring or truncation mechanism.

Examples:

- A central bank intervenes to stop an exchange rate falling below or going above certain levels.
- Dividends paid by a company may remain zero until earnings reach some threshold value.
- A government imposes price controls on some goods.
- A survey of only working women, ignoring non-working women.

In these situations, the observed data consists of a combination of measurements of some underlying *latent variable* and observations that arise when the censoring/truncation mechanism is applied.

2


Censored and Truncated Data: Definitions

- Y is *censored* when we observe X for all observations, but we only know the true value of Y for a restricted range of observations. Values of Y in a certain range are reported as a single value or there is significant clustering around a value, say 0.
 - If $Y = k$ or $Y > k$ for all Y , then Y is *censored from below* or *left-censored*.
 - If $Y = k$ or $Y < k$ for all Y , then Y is *censored from above* or *right-censored*.

We usually think of an uncensored Y , Y^* , the true value of Y when the censoring mechanism is not applied. We typically have all the observations for $\{Y, X\}$, but not $\{Y^*, X\}$.


- Y is *truncated* when we only observe X for observations where Y would not be censored. We do not have a full sample for $\{Y, X\}$, we exclude observations based on characteristics of Y .

Censored and Truncated Data: Example 1



- We observe the full range of Y and the full range of X .


Censored and Truncated Data: Example 2



- If $Y \geq 6$, we do not know its exact value.


Example: A Central Bank intervenes if the exchange rate hits the band's upper limit.
 \Rightarrow If $S_t \geq \bar{E} \Rightarrow S_t = \bar{E}$

Censored and Truncated Data: Example 2



- The pdf of the exchange rate, S_t , is a mixture of discrete (mass at $S_t = \bar{E}$) and continuous (Prob[$S_t < \bar{E}$]) distributions.


Censored and Truncated Data: Example 3



- If $Y \leq 5$, we do not know its exact value.


Example: A Central Bank intervenes if the exchange rate hits the band's lower limit. \Rightarrow If $S_t \leq \bar{E} \Rightarrow S_t = \bar{E}$.

Censored and Truncated Data: Example 3




- The pdf of the observable variable, y , is a mixture of discrete (prob. mass at $Y=5$) and continuous ($\text{Prob}[Y^* > 5]$) distributions.

Censored and Truncated Data: Example 3



- Under censoring we assign the full probability in the censored region to the censoring point, 5.


Censored and Truncated Data: Example 4



- If $Y < 3$, the value of X (or Y) is unknown. (*Truncation from below.*)

Example: If a family's income is below certain level, we have no information about the family's characteristics.

Censored and Truncated Data: Example 4



- Under data censoring, the censored distribution is a combination of a pmf plus a pdf. They add up to 1. We have a different situation under truncation. To create a pdf for Y we will use a conditional pdf.

Censored Normal

- Moments:

Let $y^* \sim N(\mu^*, \sigma^2)$ and $\alpha = (c - \mu^*)/\sigma$. Then

- Data:

$$y_i = \begin{cases} y_i^* & \text{if } y_i^* \geq c \\ c & \text{if } y_i^* \leq c \end{cases}$$

- $\text{Prob}(y = c | x) = \text{Prob}(y^* \leq c | x) = \text{Prob}[(y^* - \mu^*)/\sigma \leq (c - \mu^*)/\sigma | x]$
 $= \text{Prob}[z \leq (c - \mu^*)/\sigma | x] = \Phi(\alpha)$

- $\text{Prob}(y > c | x) = \text{Prob}(y^* > c | x) = 1 - \Phi(\alpha)$

- First Moment

- Conditional: $E[y | y^* > c] = \mu^* + \sigma \lambda(\alpha)$

- Unconditional: $E[y] = \Phi(\alpha) c + (1 - \Phi(\alpha)) [\mu^* + \sigma \lambda(\alpha)]$

Censored Normal

- To get the first moment, we use a useful result –proven later- for a truncated normal distribution. If v is a standard normal variable and the truncation is from below at c , a constant, then

$$E(v | v > c) = F^{-1} f = \frac{\phi(c)}{1 - \Phi(c)}$$

- In our conditional model, $c = -(\mathbf{x}_i' \boldsymbol{\beta})$. (Note that the expectation is also conditioned on \mathbf{x} , thus \mathbf{x} is treated as a constant.)

- Note: The ratio $F_i^{-1} f_i$ (a pdf divided by a CDF) is called *Inverse Mill's ratio*, usually denoted by $\lambda(\cdot)$ –the *hazard function*. If the truncation is from above:

$$\lambda(c) = -\frac{\phi(c)}{1 - \Phi(c)}$$

13

Censored Normal


- To get the first moment, we use a useful result –proven later- for a truncated normal distribution. If v is a standard normal variable and the truncation is from below at c , a constant, then

$$E(v | v > c) = F^{-1} f = \frac{\phi(c)}{1 - \Phi(c)} = \frac{\phi(-c)}{\Phi(-c)}$$

- In our conditional model, $c = -(\mathbf{x}_i' \boldsymbol{\beta})$. (Note that the expectation is also conditioned on \mathbf{x} , thus \mathbf{x} is treated as a constant.)

- Note: The ratio $F_i^{-1} f_i$ (a pdf divided by a CDF) is called *Inverse Mill's ratio*, usually denoted by $\lambda(\cdot)$ –the *hazard function*. If the truncation is from above:

$$\lambda(c) = -\frac{\phi(c)}{1 - \Phi(c)}$$



14

Censored Normal

- Moments (continuation):

- Unconditional first moment:

$$E[y] = \Phi(\alpha) c + (1 - \Phi(\alpha)) [\mu^* + \sigma \lambda(\alpha)]$$

If $c=0 \Rightarrow E[y] = (1 - \Phi(-\mu^*/\sigma)) [\mu^* + \sigma \lambda(\alpha)] = \Phi(\mu^*/\sigma) [\mu^* + \sigma \lambda(\alpha)]$

- Second Moment

$$\text{Var}(y) = \sigma^2 (1 - \Phi(\alpha)) \left[\begin{array}{l} (1 - \delta) \\ + (\alpha - \lambda)^2 \Phi(\alpha) \end{array} \right]$$

where $\delta = \lambda^2 - \lambda \alpha$ $0 \leq \delta \leq 1$


15

Censored Normal – Hazard Function

- The moments depend on $\lambda(c)$, the Inverse Mill's ratio or hazard rate evaluated at c :

$$\lambda(c) = \frac{\phi(c)}{1 - \Phi(c)} = \frac{\phi(-c)}{\Phi(-c)}$$

It is a monotonic function that begins at zero (when $c = -\infty$) and asymptotes at infinity (when $c = \infty$). See plot below for $(-c)$.



16

Truncated Normal

- Moments:

Let $y^* \sim N(\mu^*, \sigma^2)$ and $\alpha = (c - \mu^*)/\sigma$.

- First moment:

$$E[y^* | y > c] = \mu^* + \sigma \lambda(\alpha) \quad \leftarrow \text{This is the truncated regression.}$$

=> If $\mu > 0$ and the truncation is from below –i.e., $\lambda(\alpha) > 0$ –, the mean of the truncated variable is greater than the original mean

Note: For the standard normal distribution $\lambda(\alpha)$ is the mean of the truncated distribution.

- Second moment:

$$\text{Var}[y^* | y > c] = \sigma^2 [1 - \delta(\alpha)] \quad \text{where } \delta(\alpha) = \lambda(\alpha) [\lambda(\alpha) - \alpha]$$


=> Truncation reduces variance! This result is general, it applies to upper or lower truncation given that $0 \leq \delta(\alpha) \leq 1$

17

Truncated Normal

Model: $y_i^* = X_i\beta + \varepsilon_i$

Data: $y = y^* | y^* > 0$



- Truncated regression model:

$$E(y_i | y_i^* > 0, X_i) = X_i\beta + \sigma\lambda_i$$

18

Censored and Truncated Data: Intuition

- To model the relation between the observed y and x , we consider a latent variable y^* that is subject to censoring/truncation. A change in x affects y only through the effect of x on y^* .

- Model the true, latent variable: $y^* = f(x) = \mathbf{X} \boldsymbol{\beta} + \text{error}$

- Observed Variable: $y = h(y^*)$

- Q: What is this latent variable?

Considered the variable “Dividends paid last quarter”?

- For all the respondents with $y=0$ (*left censored from below*), we think of a latent variable for “excess cash” that underlies “dividends paid last quarter.” Extremely cash poor would pay negative dividends if that were possible.

Censored and Truncated Data: Problems

- In the presence of censoring/truncation, we have a dependent variable, y , with special attribute(s):

- (1) constrained and

- (2) clustered of observations at the constraint.

Examples:

- Consumption (1, not 2)

- Wage changes (2, not 1)

- Labor supply (1 & 2)

- These attributes create problems for the linear model.

Censored and Truncated Data: Problems

- Censoring in a regression framework (from Ruud).




Figure 28.2 Censored regression.

- If y is constrained and if there is clustering
 - OLS on the complete sample biased and inconsistent.
 - OLS on the unclustered part biased and inconsistent.

Data Censoring and Corner Solutions

- The dependent variable is partly continuous, but with positive mass at one or more points.

Two different cases:

(1) Data censoring (from above or below)

- Surveys for wealth (upper category \$250,000 or more)
- Duration in unemployment
- Demand for stadium tickets

=> Y^* is not observed because of constraints, selection technique or measuring method (data problem)

- We are interested in the effect of x on y^* : $E(y^* | \mathbf{x})$.

=> If $y_i^* = x_i' \beta + \epsilon_i$, β_j measures the effect of a change of x_j on y^* .

Data Censoring and Corner Solutions

- (2) Corner solutions: A significant fraction the data has zero value.
- Hours worked by married women: Many married women do not work –i.e, zero worked hours are reported.
 - Luxury goods, charitable donations, alcohol consumption, etc.
- => Y^* cannot be observed because of the nature of topic.
- Contrast to data censoring: Observing the dependent variable is not a problem (For modeling, we will use a latent model, y^* .)
- We are interested in $E(y | \mathbf{x})$, we want to study the effect of a change in education on y , hours worked by married women.
- => If $y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i$, β_j is not what we are interested.

Corner Solutions

- These data sets are typical of microeconomics. We think of these data sets as a result of utility maximization problems with two decisions:
- (1) Do something or not (work or not, buy IBM or not, donate or not)
- => This is binary choice problem. The *participation* decision ($y=0$ or $y>0$) part.
- (2) How much we do something (how many hours married women work, how much to invest on IBM, how much to donate to charity).
- => This a truncated sample problem. The *amount* decision (how much y is if it is positive) part.

Tobit Model (Censored Normal Regression)

- Example: We are interested on the effect of education, x , on the married women's hours worked, y .
- A model for the latent variable y^* , which is only partially observed:

$$y_i^* = \beta_0 + \beta_1 x_i + \varepsilon_i = \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i, \quad \varepsilon_i \sim N(0, \sigma^2)$$
- Data (truncated sample)
 - If $y_i^* > 0 \Rightarrow y_i = \text{Actual hours worked} = y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i$.
 - If $y_i^* \leq 0 \Rightarrow y_i = 0$ (y^* can be negative, but if it is, $y=0$)
- Probability Model -- $\varepsilon_i \sim N(0, \sigma^2)$
 - $\text{Prob}(y=0 | x) = \text{Prob}(y^* \leq 0 | x) = \text{Prob}[(y^* - \mathbf{X}\boldsymbol{\beta})/\sigma \leq (0 - \mathbf{X}\boldsymbol{\beta})/\sigma | x]$
 $= \text{Prob}[z \leq -\mathbf{X}\boldsymbol{\beta}/\sigma | x] = \Phi(-\mathbf{X}\boldsymbol{\beta}/\sigma) = 1 - \Phi(\mathbf{X}\boldsymbol{\beta}/\sigma)$
 - $\text{Prob}(y>0 | x) = \text{Prob}(y^* > 0 | x) = 1 - \Phi(-\mathbf{X}\boldsymbol{\beta}/\sigma) = \Phi(\mathbf{X}\boldsymbol{\beta}/\sigma)$ 25


Tobit Model (Censored Normal Regression)

- Expectations of interest:
 - Unconditional Expectation

$$\begin{aligned} E[y | x] &= \text{Prob}(y>0 | x) * E[y | y>0, x] + \text{Prob}(y=0 | x) * 0 \\ &= \text{Prob}(y>0 | x) * E[y | y>0, x] \\ &= \Phi(\mathbf{X}\boldsymbol{\beta}/\sigma) * E[y^* | y>0, x] \end{aligned}$$
 - Conditional Expectation (Recall: $E[y | y^* > c, x] = \mu^* + \sigma \lambda(\alpha)$)

$$E[y | y>0, x] = \mathbf{x}_i' \boldsymbol{\beta} + \sigma \lambda(\mathbf{x}_i' \boldsymbol{\beta}/\sigma)$$
- Remark: The presented Tobit model –also called *Type I Tobit Model*– can be written as a combination of two models:
- (1) A Probit model: It determines whether $y=0$ (No) or $y>0$ (Yes).
 - (2) A truncated regression model for $y>0$. 26

Tobit Model – Sample Data



- Expectations of interest:
 $E[y | x]$, $E[y | y > 0, x]$.
 $E[y^* | x]$

27

Tobit Model - OLS

- Suppose we do OLS, only for the part of the sample with $y > 0$. The estimated model:

$$y_i = \mathbf{x}_i' \boldsymbol{\beta} + v_i, \quad \text{for } y_i > 0$$

- Let's look at the density of $v_i, f_v(\cdot)$, which must integrate to 1:

$$\int_{-x_i' \boldsymbol{\beta}}^{\infty} f_v(\eta) d\eta = 1$$

- The ϵ_i 's density, normal by assumption in the Tobit Model:

$$\int_{-x_i' \boldsymbol{\beta}}^{\infty} f_\epsilon(\eta) d\eta = F_i = \int_{-\infty}^{x_i' \boldsymbol{\beta}} f_\epsilon(\eta) d\eta = \int_{-\infty}^{x_i' \boldsymbol{\beta}} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{\eta}{\sigma}\right)^2}$$

- Then, $f_v(\cdot)$ can be written as:

$$f_v = F_i^{-1} f_\epsilon = F_i^{-1} \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{v_i}{\sigma}\right)^2}$$

28

Tobit Model - OLS

- The pdf of v_i and the pdf of ε_i are different. In particular, The ε_i 's density, normal by assumption in the Tobit Model:

$$\begin{aligned}
 E[v_i] &= \int_{-x_i'\beta}^{\infty} \eta f_v(\eta) d\eta = F_i^{-1} \int_{-x_i'\beta}^{\infty} \eta f_\varepsilon(\eta) d\eta \\
 &= F_i^{-1} \int_{-x_i'\beta}^{\infty} \eta \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{\eta}{\sigma}\right)^2} d\eta = F_i^{-1} \int_{-x_i'\beta}^{\infty} \frac{\eta}{\sigma} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{\eta}{\sigma}\right)^2} d\eta \\
 &= F_i^{-1} [-\sigma f_\varepsilon(\eta)]_{-x_i'\beta}^{\infty} && \text{integration by substitution} \\
 &= \sigma F_i^{-1} f_i \neq 0 && (f_i = f_\varepsilon(x_i'\beta))
 \end{aligned}$$

- Then, $E[v_i | \mathbf{x}] = \sigma F_i^{-1} f_i = \sigma \lambda(\mathbf{x}_i'\beta) \neq 0$ (and it depends on $\mathbf{x}_i'\beta$)
 $\Rightarrow E[y_i | y_i > 0, \mathbf{x}_i'\beta] = \mathbf{x}_i'\beta + \sigma \lambda(\mathbf{x}_i'\beta)$
 \Rightarrow OLS in truncated part is *biased* (omitted variables problem).

29

Tobit Model - OLS

- OLS in truncated part only is *biased*. We have an omitted variables problem. With a bit more work, we can show OLS is inconsistent too (show it!)

- Also $E[v_i^2 | \mathbf{x}] = \sigma^2 - \sigma^2 \mathbf{x}_i'\beta \lambda(\mathbf{x}_i'\beta)$

- \Rightarrow The error term, v , not only has non-zero mean but it is also heteroskedastic.

30

Tobit Model - NLLS

- Now, we can write

$$y_i = E[y_i | y_i > 0, \mathbf{x}_i' \boldsymbol{\beta}] + \boldsymbol{\varepsilon}_i = \mathbf{x}_i' \boldsymbol{\beta} + \sigma \lambda(\mathbf{x}_i' \boldsymbol{\beta}) + \boldsymbol{\varepsilon}_i, \quad \text{for } y_i > 0$$

- There is a non-linear relation between x_i on y . NLLS is a possibility here, though we need a consistent estimator of $\boldsymbol{\beta}$ to evaluate $\lambda(\mathbf{x}_i' \boldsymbol{\beta})$ and it is not clear where to get it. Since we have heteroscedasticity, we need to allow for it. Weighted NLLS may work well.

- Note: A non-linear relation appears even if all observations are used (positive and negative values of y):

$$\begin{aligned} E[y_i | \mathbf{x}_i' \boldsymbol{\beta}] &= \text{Prob}(y_i^* > 0 | \mathbf{x}_i) * E(y_i^* | y_i^* > 0, \mathbf{x}_i) = F_i(\mathbf{x}_i' \boldsymbol{\beta} + \sigma F_i^{-1} f_i) \\ &= \Phi(\mathbf{x}_i' \boldsymbol{\beta}) \mathbf{x}_i' \boldsymbol{\beta} + \sigma \phi(\mathbf{x}_i' \boldsymbol{\beta}) \end{aligned}$$

31

Tobit Model: Estimation

- Given our assumption for $\boldsymbol{\varepsilon}_i$, we do ML estimation.
- For women who are working, we have $y_i^* > 0 \Rightarrow y_i > 0$. Then,

$$\boldsymbol{\varepsilon}_i = y_i - \beta_0 + \beta_1 x_i$$

\Rightarrow Likelihood function for a working woman is given by:

$$L_i = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{(y_i - \beta_0 - \beta_1 x_i)^2}{2\sigma^2}} = \frac{1}{\sigma} \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2} \frac{(y_i - \beta_0 - \beta_1 x_i)^2}{\sigma^2}} = \frac{1}{\sigma} \phi\left(\frac{y_i - \beta_0 - \beta_1 x_i}{\sigma}\right)$$

- For women who are not working, we have $y_i^* \leq 0 \Rightarrow y_i = 0$. The likelihood contribution is the probability that $y_i^* \leq 0$, which is given by

$$\begin{aligned} L_i &= P(y_i^* \leq 0) = P(\beta_0 + \beta_1 x_i + \boldsymbol{\varepsilon}_i \leq 0) \\ &= P(\boldsymbol{\varepsilon}_i \leq -(\beta_0 + \beta_1 x_i)) = P\left(\frac{\boldsymbol{\varepsilon}_i}{\sigma} \leq -\frac{\beta_0 + \beta_1 x_i}{\sigma}\right) \\ &= \Phi\left(-\frac{\beta_0 + \beta_1 x_i}{\sigma}\right) = 1 - \Phi\left(\frac{\beta_0 + \beta_1 x_i}{\sigma}\right) \end{aligned}$$

32

Tobit Model: Estimation

- To summarize,

$$L_i = \frac{1}{\sigma} \phi\left(\frac{y_i - \beta_0 - \beta_1 x_i}{\sigma}\right) \quad \text{if } y_i^* > 0$$

$$= 1 - \Phi\left(\frac{\beta_0 + \beta_1 x_i}{\sigma}\right) \quad \text{if } y_i^* \leq 0$$

- We have a combination of a pdf (for the observed part of the distribution) and a CDF (for the truncated part of the distribution): a linear part and a Probit part.
- Let D_i be a dummy variable that takes 1 if $y_i > 0$. Then, the above likelihood for consumer i can be written as.

$$L_i = \left[\frac{1}{\sigma} \phi\left(\frac{y_i - \beta_0 - \beta_1 x_i}{\sigma}\right) \right]^{D_i} \left[1 - \Phi\left(\frac{\beta_0 + \beta_1 x_i}{\sigma}\right) \right]^{1-D_i}$$

33

Tobit Model: Estimation

- The likelihood function, L , for the whole sample is:

$$L(\beta_0, \beta_1, \sigma) = \prod_{i=1}^n L_i = \prod_{i=1}^n \left[\frac{1}{\sigma} \phi\left(\frac{y_i - \beta_0 - \beta_1 x_i}{\sigma}\right) \right]^{D_i} \left[1 - \Phi\left(\frac{\beta_0 + \beta_1 x_i}{\sigma}\right) \right]^{1-D_i}$$

- The values of β_0, β_1 and σ that maximize the likelihood function are the Tobit estimators of the parameters.
- As usual, we work with the $\text{Log}(L)$:

$$L(\beta_0, \beta_1, \sigma) = \sum_{i=1}^n D_i \log \left[\frac{1}{\sigma} \phi\left(\frac{y_i - \beta_0 - \beta_1 x_i}{\sigma}\right) \right] + (1 - D_i) \log \left[1 - \Phi\left(\frac{\beta_0 + \beta_1 x_i}{\sigma}\right) \right]$$

$$= \frac{N}{2} [\log(\sigma^2) + \log(2\pi)] +$$

$$+ \sum_{i=1}^n D_i \left[-\frac{(y_i - \beta_0 - \beta_1 x_i)^2}{2\sigma^2} + (1 - D_i) \log \left[1 - \Phi\left(\frac{\beta_0 + \beta_1 x_i}{\sigma}\right) \right] \right]$$

34

Tobit Model: Estimation – Information Matrix

- Amemiya (1973) presents the following representation for the information matrix:

$$I(\Theta)_{((K+1) \times (K+1))} = \begin{bmatrix} \sum_{i=1}^T a_i X_i X_i' & \sum_{i=1}^T b_i X_i \\ \sum_{i=1}^T b_i X_i' & \sum_{i=1}^T c_i \end{bmatrix}$$

$(K \times K)$
 $(K \times 1)$
 $(1 \times K)$
 (1×1)

where:

$$z_i = \frac{X_i' \beta}{\sigma} \quad a_i = \frac{-1}{\sigma^2} \left(z_i f(z_i) - \frac{f(z_i)^2}{1 - F(z_i)} - F(z_i) \right)$$


$$b_i = \frac{1}{2\sigma^3} \left(z_i^2 f(z_i) + f(z_i) - \frac{z_i f(z_i)^2}{1 - F(z_i)} \right)$$

$$c_i = -\frac{1}{4\sigma^4} \left(z_i^3 f(z_i) + z_i f(z_i) - \frac{z_i^2 f(z_i)^2}{1 - F(z_i)} - 2F(z_i) \right)$$

35

Tobit Model: Application I – Female Labor Supply

- **Corner Solution Case – DATA:** Female labor supply (mroz.wf1)



Variable	Obs	Mean	Std. Dev.	Min	Max
hours	753	740.5764	871.3142	0	4950
nwifeinc	753	20.12896	11.6348	-.0290575	96
educ	753	12.28685	2.280246	5	17
exper	753	10.63081	8.06913	0	45
age	753	42.53785	8.072574	30	60
kidsge6	753	1.353254	1.319874	0	8
kidslt6	753	.2377158	.523959	0	3

Tobit Model : Application I – OLS (all y)

- OLS whole sample (N=753)

Dependent Variable: HOURS

Method: Least Squares

Included observations: 753

White Heteroskedasticity-Consistent Standard Errors & Covariance

Variable	Coefficient	Std. Error	t-Statistic	Prob.
NWIFEINC	-3.446636	2.240662	-1.538222	0.1244
EDUC	28.76112	13.03905	2.205768	0.0277
EXPER	65.67251	10.79419	6.084062	0.0000
EXPER^2	-0.700494	0.372013	-1.882983	0.0601
AGE	-30.51163	4.244791	-7.188018	0.0000
KIDSLT6	-442.0899	57.46384	-7.693359	0.0000
KIDSGE6	-32.77923	22.80238	-1.437535	0.1510
C	1330.482	274.8776	4.840273	0.0000

Tobit Model : Application I – OLS (y>0)

- OLS subsample - hours>0 (N=428)

Dependent Variable: HOURS

Method: Least Squares

Sample: 1 753 IF HOURS>0

Included observations: 428

White Heteroskedasticity-Consistent Standard Errors & Covariance

Variable	Coefficient	Std. Error	t-Statistic	Prob.
NWIFEINC	0.443851	3.108704	0.142777	0.8865
EDUC	-22.78841	16.09281	-1.416061	0.1575
EXPER	47.00509	15.38725	3.054808	0.0024
EXPER^2	-0.513644	0.417384	-1.230627	0.2192
AGE	-19.66352	5.845279	-3.364001	0.0008
KIDSLT6	-305.7209	125.4802	-2.436407	0.0152
KIDSGE6	-72.36673	31.28480	-2.313159	0.0212
C	2056.643	351.4502	5.851875	0.0000

Tobit Model : Application I – Tobit

- Tobit whole sample (N=753)

Dependent Variable: HOURS
 Method: ML - Censored Normal (TOBIT) (Quadratic hill climbing)
 Included observations: 753
 Left censoring (value) at zero

	Coefficient	Std. Error	z-Statistic	Prob.
NWIFEINC	-8.814243	4.459100	-1.976687	0.0481
EDUC	80.64561	21.58324	3.736493	0.0002
EXPER	131.5643	17.27939	7.613943	0.0000
EXPER^2	-1.864158	0.537662	-3.467155	0.0005
AGE	-54.40501	7.418502	-7.333693	0.0000
KIDSLT6	-894.0217	111.8780	-7.991039	0.0000
KIDSGE6	-16.21800	38.64139	-0.419705	0.6747
C	965.3053	446.4361	2.162247	0.0306
Error Distribution				
SCALE:C(9)	1122.022	41.57910	26.98523	0.0000
Left censored obs	325	Right censored obs	0	
Uncensored obs	428	Total obs	753	

Partial Effects (marginal effects)

- The estimated parameters β_j measures the effect of x_j on y^* . But in corner solutions, we are interested in the effect of x_j on actual y .

- We calculate partial effects based on two results already derived:

- For positive y 's –i.e., $y_i > 0$:

$$E[y_i | y_i > 0, \mathbf{x}_i' \boldsymbol{\beta}] = \mathbf{x}_i' \boldsymbol{\beta} + \sigma \lambda(\mathbf{x}_i' \boldsymbol{\beta})$$

- For all y 's –i.e., $y_i \geq 0$:

$$E[y_i | \mathbf{x}_i' \boldsymbol{\beta}] = \text{Prob}(y^* > 0 | \mathbf{x}) * E(y^* | y > 0, \mathbf{x}) = F_1[\mathbf{x}_i' \boldsymbol{\beta} + \sigma^2 \lambda(\mathbf{x}_i' \boldsymbol{\beta})]$$

- The partial effects are given by

$$(1) \quad \frac{\partial E(y | y > 0, \mathbf{x})}{\partial x_k} = \beta_k \left\{ 1 - \lambda\left(\frac{\mathbf{x}'\boldsymbol{\beta}}{\sigma}\right) \left[\frac{\mathbf{x}'\boldsymbol{\beta}}{\sigma} + \lambda\left(\frac{\mathbf{x}'\boldsymbol{\beta}}{\sigma}\right) \right] \right\} = \beta_k (1 - \delta_k)$$

(1) measures the effect of an x_k change on y for working women⁴⁰

Partial Effects (marginal effects)

=> β_k overstates the marginal impact of a change in X_k

$$2. \frac{\partial E(y|x)}{\partial x_k} = \beta_k \Phi\left(\frac{x'\beta}{\sigma}\right)$$

(2) measures the overall effect of an x_k change on hours worked.

- Both partial effects depend on \mathbf{x} . Therefore, they vary by person.
- We are interested in the overall effect rather than the effect for a specific person in the data. Two ways to do this computation:
 - At the sample average: Plug the mean of \mathbf{x} in the above formula.
 - Average of partial effects: Compute the partial effect for each individual in the data. Then, compute the average.

41

Partial Effects – Application I ($y > 0$)

```
. tobit hours nwifeinc educ exper expersq age kidslt6 kidsge6, ll(0)
Tobit regression              Number of obs =      753
                             LR chi2(7)         =    271.59
                             Prob > chi2        =    0.0000
                             Pseudo R2         =    0.0343
Log likelihood = -3819.0946
```

hours	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
nwifeinc	-8.814243	4.459096	-1.98	0.048	-17.56811	-.0603724
educ	80.64561	21.58322	3.74	0.000	38.27453	123.0167
exper	131.5643	17.27938	7.61	0.000	97.64231	165.4863
expersq	-1.864158	.5376615	-3.47	0.001	-2.919667	-.8086479
age	-54.40501	7.418496	-7.33	0.000	-68.96862	-39.8414
kidslt6	-894.0217	111.8779	-7.99	0.000	-1113.655	-674.3887
kidsge6	-16.218	38.64136	-0.42	0.675	-92.07675	59.64075
_cons	965.3053	446.4358	2.16	0.031	88.88528	1841.725
/sigma	1122.022	41.57903			1040.396	1203.647

```
obs. summary:      325 left-censored observations at hours=0
                  428 uncensored observations
                  0 right-censored observations
```

```
*****
*Compute the Partial effect *
*at average of educ *
*on hours for working women *
*manually *
*****
. predict xbeta, xb
. egen avxbeta=mean(xbeta)
. gen avxbsig=avxbeta/_b[sigma]
. gen lambda=normalden(avxbsig)/normal(avxbsig)
. gen partial=_b[educ]*(1-lambda*(avxbsig+lambda))
. su partial
```

Variable	Obs	Mean	Std. Dev.	Min	Max
partial	753	34.27517	0	34.27517	34.27517

Partial effect at average
for *working women*:
Computing manually.

42

Partial Effects – Application I (y>0)

```
. tobit hours mwifeinc educ exper expersq age kids1t6 kidsge6, ll(0)
```

```
Tobit regression      Number of obs   =    753
                     LR chi2(7)      =   271.59
                     Prob > chi2     =    0.0000
Log likelihood = -3819.0946      Pseudo R2       =    0.0343
```

hours	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]
mwifeinc	-8.814243	4.459096	-1.98	0.048	-17.56811 -0.0603724
educ	80.64561	21.58322	3.74	0.000	38.27453 123.0167
exper	131.5643	17.27938	7.61	0.000	97.64231 165.4863
expersq	-1.864158	.5376615	-3.47	0.001	-2.919667 -0.8086479
age	-54.40301	7.418496	-7.33	0.000	-68.96862 -39.8414
kids1t6	-894.0217	111.8779	-7.99	0.000	-1113.655 -674.3887
kidsge6	-16.218	38.64136	-0.42	0.675	-92.07675 59.64075
_cons	965.3053	446.4358	2.16	0.031	88.88528 1841.725
/sigma	1122.022	41.57903			1040.396 1203.647

```
obs. summary:      325 left-censored observations at hours<=0
                   428 uncensored observations
                   0 right-censored observations
```

```
*****
* Compute the partial effect *
* at average of educ on hours *
* for working women automatically *
*****
. mfx, predict(e(0,.)) varlist(educ)
```

```
Marginal effects after tobit
y = E(hours|hours>0) (predict, e(0,.))
= 1012.0327
```

variable	dy/dx	Std. Err.	z	P> z	[95% C.I.]	x
educ	34.27517	9.11708	3.76	0.000	16.406 52.1443	12.2869

Partial effect at average for *working women*: Compute automatically.

Partial Effects – Application I (all y)

```
*****
* Compute the Partial effect at average *
* of education for the entire observation*
* manually *
*****
```

```
. gen partial_all=b[educ]*normal(avxbsig)
```

```
. su partial_all
```

Variable	Obs	Mean	Std. Dev.	Min	Max
partial_all	753	48.73409	0	48.73409	48.73409

```
*****
* Compute the Partial effect at average *
* of education for the entire observation*
* automatically *
*****
```

```
. mfx, predict(y*(0,.)) varlist(educ)
```

```
Marginal effects after tobit
y = E(hours*|hours>0) (predict, y*(0,.))
= 611.57078
```

variable	dy/dx	Std. Err.	z	P> z	[95% C.I.]	x
educ	48.73409	12.963	3.76	0.000	23.3263 74.1419	12.2869

Partial effect at average for *all observations*: Compute manually.

Partial effect at average for *all observations*: Compute automatically.

Partial Effects – Application I (all y)

- Now, we can compare the marginal effect of education on actual hours worked.
- We compare OLS (whole sample) and Tobit estimates, on the basis of the marginal effect of education actual y, *for an average individual*.

OLS	TOBIT	
$\hat{\beta}_{k,OLS}$	$\beta_k \Phi \left(\frac{\bar{x}'\beta}{\sigma} \right)$	→ OLS <u>underestimates</u>
28.76	$\begin{matrix} \uparrow \\ 80.65 \quad 0.604 \\ \hline 48.73 \end{matrix}$	the effect of education on the labor supply (in the average of the explanatory variables).

Interpretation: On average, an additional year of education increases the labor supply by 48,7 hours (for an average individual).

Tobit Model: Heteroscedasticity

- In a regression model, we scale the observations by their standard deviation (x_i/σ_i) transforming the model back to CLM framework
- In the Tobit model, we naturally work with the likelihood. The Log L for the homoscedastic Tobit model is:

$$L(\beta_0, \beta_1, \sigma) = \frac{N}{2} [\log(\sigma^2) + \log(2\pi)] + \sum_{i=1}^n \left[-D_i \frac{(y_i - \beta_0 - \beta_1 x_i)^2}{2\sigma^2} + (1 - D_i) \log \left[1 - \Phi \left(\frac{\beta_0 + \beta_1 x_i}{\sigma} \right) \right] \right]$$

- Introducing heteroscedasticity in the Log L:

$$L(\beta_0, \beta_1, \sigma_1, \sigma_2, \dots, \sigma_N) = \frac{N}{2} \log(2\pi) + \sum_{i=1}^n \left[D_i [\log(\sigma_i) - \frac{(y_i - \beta_0 - \beta_1 x_i)^2}{2\sigma_i^2}] + (1 - D_i) \log \left[1 - \Phi \left(\frac{\beta_0 + \beta_1 x_i}{\sigma_i} \right) \right] \right]_{46}$$

Tobit Model: Heteroscedasticity

- Now, we went from $k+1$ parameters to $k+N$ parameters. Impossible to estimate with N observations.
- Usual solution: Model heteroscedasticity, dependent on a few parameters: $\sigma_i^2 = \sigma_i(\alpha)$.

Example: Exponential: $\sigma_i^2 = \exp(\mathbf{z}_i' \alpha)$. Then,

$$L(\beta_0, \beta_1, \alpha) = \frac{N}{2} \log(2\pi) + \sum_{i=1}^n \left[D_i \left[\mathbf{z}_i' \alpha - \frac{(y_i - \beta_0 - \beta_1 x_i)^2}{2 \exp(\mathbf{z}_i' \alpha)} \right] + (1 - D_i) \log \left[1 - \Phi \left(\frac{\beta_0 + \beta_1 x_i}{\sqrt{\exp(\mathbf{z}_i' \alpha)}} \right) \right] \right]$$

- The marginal effects get more complicated under the heteroscedastic Tobit model: an exogenous variable, say income, could impact both numerator and denominator of the standardization ratio, $X_i \beta / \sigma_i$ 47

Heteroscedasticity – Partial Effects

- Partial effects get more complicated under the heteroscedastic Tobit model: an exogenous variable, say income, could impact both numerator, $X_i \beta$, and denominator σ_i . Ambiguous signs are possible.
- Suppose we have w_j affecting both \mathbf{x}_i and \mathbf{z}_i . Then,

$$r_i = \frac{x_i' \beta_i}{\sigma_i} = \frac{x_i' \beta_i}{\exp(\mathbf{z}_i' \alpha / 2)}$$

$$\frac{\partial r_i}{\partial w_{ji}} = \frac{\beta_i \sigma_i - (x_i' \beta_i)(\alpha_i / 2) \sigma_i}{\sigma_i^2} = \frac{\beta_i - (x_i' \beta_i)(\alpha_i / 2)}{\sigma_i}$$

$$\frac{\partial \Phi(r_i)}{\partial w_{ji}} = \phi(r_i) \frac{\partial r_i}{\partial w_{ji}} = \phi(r_i) \frac{\beta_i - (x_i' \beta_i)(\alpha_i / 2)}{\sigma_i}$$

$$\frac{\partial E[y_i | y_i > 0]}{\partial w_{ji}} = \beta_i - \sigma_i \frac{\partial r_i}{\partial w_{ji}} \left(r_i \frac{\phi(r_i)}{\Phi(r_i)} + \frac{\phi(r_i)^2}{\Phi(r_i)^2} \right)$$

Heteroscedasticity – Partial Effects - Application

- Canadian FAFH expenditures: 9,767 HH's, 21.2% with \$0 expenditures.
- Dependent variable is bi-weekly FAFH expenditures
Exogenous Variables: HHInc, Kids Present?, FullTime? Provincial Dummy Variables.
- $\sigma_i^2 = \exp(\gamma_0 + \gamma_1 \text{Income}_i + \gamma_2 \text{Fulltime}_i + \gamma_3 \text{Quebec}_i)$

Elasticity	Homo.		Hetero.	
	Value	S.E.	Value	S.E.
$\Phi(z)$	0.259	0.008	0.210	0.011
$E(y y>0)$	0.284	0.009	0.395	0.010
$E(y)$	0.544	0.017	0.606	0.020

49

Tobit Model – Type II

- Different ways of thinking about how the latent variable and the observed variable interact produce different Tobit Models.

- The Type I Tobit Model presents a simple relation:

$$\begin{aligned}
 - y_i &= 0 && \text{if } y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i \leq 0 \\
 &= y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i && \text{if } y_i^* = \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_i > 0
 \end{aligned}$$

The effect of the X 's on the probability that an observation is censored and the effect on the conditional mean of the non-censored observations are the same: $\boldsymbol{\beta}$.

- The Type II Tobit Model presents a more complex relation:

$$\begin{aligned}
 - y_i &= 0 && \text{if } y_i^* = \mathbf{x}_i' \boldsymbol{\alpha} + \varepsilon_{1,i} \leq 0, \quad \varepsilon_{1,i} \sim N(0,1) \\
 &= \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_{2,i} && \text{if } y_i^* = \mathbf{x}_i' \boldsymbol{\alpha} + \varepsilon_{1,i} > 0, \quad \varepsilon_{2,i} \sim N(0, \sigma_2^2)
 \end{aligned}$$

Now, we have different effects of the X 's.

50

Tobit Model – Type II

- The Type II Tobit Model:

$$\begin{aligned}
 - y_i &= 0 && \text{if } y_i^* = \mathbf{x}_i' \boldsymbol{\alpha} + \varepsilon_{1,i} \leq 0, \quad \varepsilon_{1,i} \sim N(0, \sigma_1^2 = 1) \\
 &= \mathbf{x}_i' \boldsymbol{\beta} + \varepsilon_{2,i} && \text{if } y_i^* = \mathbf{x}_i' \boldsymbol{\alpha} + \varepsilon_{1,i} > 0, \quad \varepsilon_{2,i} \sim N(0, \sigma_2^2)
 \end{aligned}$$

- A more flexible model. \mathbf{X} can have an effect on the decision to participate (Probit part) and a different effect on the amount decision (truncated regression).

- Type I is a special case: $\varepsilon_{2,i} = \varepsilon_{1,i}$ and $\boldsymbol{\alpha} = \boldsymbol{\beta}$.

Example: Age affects the decision to donate to charity. But it can have a different effect on the amount donated. We may find that age has a positive effect on the decision to donate, but given a positive donation, younger individuals donate more than older individuals.

51

Tobit Model – Type II

- The Tobit Model assumes a bivariate normal distribution for $(\varepsilon_{1,i}; \varepsilon_{2,i})$; with covariance given by $\sigma_{12} (= \rho \sigma_1 \sigma_2)$.

- Conditional expectation:

$$E[y | y > 0, \mathbf{x}] = \mathbf{x}_i' \boldsymbol{\beta} + \sigma_{12} \lambda(\mathbf{x}_i' \boldsymbol{\alpha})$$

- Unconditional Expectation

$$\begin{aligned}
 E[y | \mathbf{x}] &= \text{Prob}(y > 0 | \mathbf{x}) * E[y | y > 0, \mathbf{x}] + \text{Prob}(y = 0 | \mathbf{x}) * 0 \\
 &= \text{Prob}(y > 0 | \mathbf{x}) * E[y | y > 0, \mathbf{x}] \\
 &= \Phi(\mathbf{x}_i' \boldsymbol{\alpha}) * [\mathbf{x}_i' \boldsymbol{\beta} + \sigma_{12} \lambda(\mathbf{x}_i' \boldsymbol{\alpha})]
 \end{aligned}$$

Note: This model is known as the Heckman selection model, or the Type II Tobit model (Amemiya), or the probit selection model (Wooldridge).

52